

2023-24
English Medium

SSC- MATHEMATICS

**CHAPTERWISE
SOLVED PAPERS**

**Youth
Competition
Times**

STAFF SELECTION COMMISSION

MATHEMATICS

SSC

2023-24

CGL (TIER I & II), CHSL (10+2),
CPO-SI, MTS, GD, Selection Post

**CHAPTERWISE
Solved Papers**

**CHAPTERWISE &
SUB TOPICWISE**

Computer
Based Test

10150⁺
OBJECTIVE
QUESTIONS

Include Chapterwise Presentation of **353** Online Question Papers (All Sets)

Staff Selection Commission

SSC

Maths

Chapterwise Solved Papers

(Computer Based Test)

CHIEF EDITOR

A.K. Mahajan

EDITOR

Advocate Abhishek Singh

WRITERS


Ambuj Kumar, Anand Soni, Kamlesh Shrivastav
Vineet Shrivastav, Rahul Verma, Hemant Sharma

COMPUTER GRAPHICS BY

Balkrishna, Charan Singh, Vinay Sahu

EDITORIAL OFFICE

Youth Competition Times
12, Church Lane Prayagraj-211002

 **Mob. :** 9415650134

Email : yctap12@gmail.com

website : www.yctbooks.com

PUBLISHER DECLARATION

Edited and Published by A.K. Mahajan printed by Om Sai Offset, Prayagraj.
Youth Competition Times 12, Church Lane, Prayagraj

In order to publish the book, full care has been taken by the editor and the publisher,
Still your suggestions and queries are welcomed.

In case of any dispute, the judicial area will be Prayagraj.

Rs. : 695/-

INDEX

- Analysis Chart of SSC Maths Previous Year Exam Question Papers..... 6
- Trend Analysis of Previous Year Papers of SSC Exams Through Pie Chart and Bar Graph7-8

SECTION-1

■ Algebra	9-79
■ Problems based on Linear Equations	9
■ Problems based on Algebraic Identities	12
■ Problems based on Factors of Polynomials and Remainder Theorem.....	66
■ Problems based on Quadratic Equations and Nature of its Roots	69
■ Miscellaneous	71
■ Trigonometry	80-156
■ Problems based on Trigonometric Ratios	80
■ Problems based on Trigonometric Identities	94
■ Problems based on Trigonometric Functions	103
■ Problems based on Angular Values of Trigonometric Functions.....	111
■ Miscellaneous	143
■ Height and Distance	157-172
■ Geometry	173-295
■ Problems based on Lines and Angles.....	173
■ Problems based on Congruency and Similarity of Triangles.....	173
■ Problems based on Quadrilateral	220
■ Problems based on Square	224
■ Problems based on Rectangle	225
■ Problems based on Rhombus	226
■ Problems based on Parallelogram	228
■ Problems based on Trapezium.....	229
■ Problems based on Right angled Triangle	232
■ Problems based on Circle.....	240
■ Problems based on Tangent to Circle.....	270
■ Problems based on Equilateral Polygon.....	291
■ Miscellaneous	293
■ Co-ordinate Geometry	296-297
■ Mensuration 2D	298-335
■ Problems based on Triangle.....	298
■ Problems based on Quadrilateral.....	310
■ Problems based on Circle.....	311
■ Problems based on Square	319
■ Problems based on Rectangle	323
■ Hexagon	329
■ Semi circle.....	331
■ Rhombus.....	331
■ Miscellaneous	332

■ Mensuration 3D	336-371
■ Problems based on Cube	336
■ Problems based on Cuboid.....	339
■ Problems based on Cylinder	344
■ Problems based on Cone	353
■ Problems based on Sphere and Hemisphere	359
■ Problems based on Prism and Pyramid.....	366
■ Miscellaneous	369

SECTION-2

■ Number System.....	372-410
■ Problems based on Divisibility Rule	372
■ Problems based on Prime and Composite Numbers.....	394
■ Problems based on Factors of Numbers.....	394
■ Problems based on Unit Digit of Numbers	396
■ Problems based on Remainder Theorem.....	397
■ Problems based on Progression	403
■ Problems based on Arithmetic and Geometric Progression.....	404
■ Miscellaneous	406
■ Decimal and Fraction	411-417
■ Problems based on finding smallest and largest fractions	411
■ Problems based on Simplification of fractions.....	412
■ Miscellaneous	416
■ Indices & Surds	418-428
■ Problems based on Square Root of Numbers	418
■ Problems based on Exponents	419
■ Problems based on Surds.....	421
■ Miscellaneous	428
■ LCM and HCF.....	429-440
■ Problems based on L.C.M.....	429
■ Problems based on H.C.F.....	433
■ Mixed Problems based on H.C.F. and L.C.M.....	435
■ Miscellaneous	438
■ Simplification	441-480
■ Problem based on BODMAS Rule	441
■ Miscellaneous	475
■ Average.....	481-514
■ Simple Problems based on Average	481
■ Problems based on Average of Consecutive Numbers	497
■ Problems based on Finding Average Age/Weight	501
■ Problems based on Marks Obtained by Students in an Examination.....	507
■ Problems based on Runs Scored in Cricket Matches	510
■ Miscellaneous	512

■ Ratio and Proportion	515-543
■ Problems based on Basic Interpretation of Ratio and Proportion	515
■ Problems based on Finding New Proportion due to Increase or Decrease in Original Ratio/ Proportion	526
■ Problems based on Income and Expenditures	532
■ Problems based on Ratio of Coins and Rupees	537
■ Miscellaneous	539
■ Percentage.....	544-585
■ Problems based on Concepts of Percentage.....	544
■ Problems based on Percentage Change.....	562
■ Problems based on Percentage Change in Area and Volume.....	566
■ Problems based on Population.....	569
■ Problems based on Election.....	570
■ Problems based on Income, Expenditure and Savings.....	572
■ Problems based on Pass/Fail Candidates in an Examination.....	579
■ Miscellaneous	581
■ Profit and Loss.....	586-634
■ Problems based on Profit and Loss.....	586
■ Problems based on Finding Cost Price.....	604
■ Problems based on Finding Selling Price.....	611
■ Problems based on Buying and Selling of two items	621
■ Problems based on Buying and Selling of an Object at Special rate.....	626
■ Miscellaneous	629
■ Discount.....	635-663
■ Problems based on Discount.....	635
■ Problems based on Successive Discount.....	642
■ Problems based on Marked Price	647
■ Problems based on Selling Price.....	654
■ Miscellaneous	656
■ Simple Interest.....	664-681
■ Problems based on Fundamental of Simple Interest	664
■ If an amount becomes n times in t years at the rate of Simple Interest.....	674
■ If an amount becomes P_1 in t_1 years and P_2 in t_2 years at the fixed rate of Simple Interest.....	675
■ Miscellaneous	677
■ Compound Interest.....	682-714
■ Problems based on Fundamental formula of Compound Interest	682
■ If an amount becomes n times in t years at the rate of Compound Interest.....	696
■ Problems based on difference in Simple Interest and Compound Interest	696
■ Mixed problems on Simple and Compound Interest.....	700
■ Miscellaneous	709

■ Partnership	715-722
■ Problems based on finding the share of one person in a partnership of two or three persons	715
■ Problems based on Profit accumulated in Partitions in a partnership	718
■ Problems based on Principal Amount	721
■ Miscellaneous	722
■ Alligation	723-729
■ Time and Work.....	730-776
■ Problems based on Time and Work.....	730
■ Problems based on Remaining Work	748
■ When a person leaves the work incomplete during the process	759
■ When Additional Persons are hired to complete the work after commencement of work	763
■ Problems based on Efficiency.....	764
■ Problems based on Work and Wages	772
■ Miscellaneous	776
■ Pipes and Cisterns	777-789
■ Problems based on Tap and Tank	777
■ When one tap performs the role of filling while the other one performs the role of emptying.....	781
■ When in between the tap is opened or closed.....	786
■ When the Taps are opened alternatively	788
■ Miscellaneous	788
■ Speed, Time and Distance.....	790-817
■ Problems based on Speed, Time and Distance	790
■ When Distance is Constant	802
■ When Time is Constant	809
■ Problems based on Average Speed.....	810
■ Miscellaneous	811
■ Problems Related to Train.....	818-831
■ Simple Problem related to Train.....	818
■ When the Train/Person Moves in Opposite Direction of other Train/Person.....	821
■ When the train/person moves in same direction of other train/ person.....	825
■ Miscellaneous	827
■ Boat and Stream	832-838
■ Problems based on Speed of Boat or Swimmer.....	832
■ Problems based on Speed of Stream	835
■ Problems based on Time	836
■ Miscellaneous	837
■ Age Problems	839-845
■ Data Interpretation	846-912
■ Problems based on Bar-Graph	846
■ Problems based on Tables.....	867
■ Problems based on Pie-Chart.....	888
■ Problems based on Line Diagram.....	909
■ Problems based on Histogram	912



Analysis Chart of Question Papers of Various Previous Exam of SSC

Sr. No.	Exam	Exam Year	Total Questions	Total Question of Maths
1.	SSC CGL	2022	21	$21 \times 25 = 525$
2.	SSC CHSL	2022	42	$42 \times 25 = 1050$
3.	SSC MTS	2021	42	$42 \times 25 = 1050$
4.	SSC CPO-SI	2020	6	$6 \times 50 = 300$
5.	SSC Selection Post Phase VIII (Graduate Level)	2020	4	$4 \times 25 = 100$
6.	SSC Selection Post Phase VIII (H.S. Level)	2020	3	$3 \times 25 = 75$
7.	SSC Selection Post Phase VIII (Matriculation Level)	2020	5	$5 \times 25 = 125$
8.	SSC CGL (Tier-II)	2020	3	$3 \times 100 = 300$
9.	SSC CHSL	2020	36	$36 \times 25 = 900$
10.	SSC CGL (Tier-I)	2020	18	$18 \times 25 = 450$
11.	SSC CPO-SI	2019	8	$8 \times 50 = 400$
12.	SSC Selection Post Phase VII (Graduate Level)	2019	4	$4 \times 25 = 100$
13.	SSC Selection Post Phase VII (H.S. Level)	2019	4	$4 \times 25 = 100$
14.	SSC Selection Post Phase VII (Matriculation Level)	2019	4	$4 \times 25 = 100$
15.	SSC CGL (Tier-II)	2019	3	$3 \times 100 = 300$
16.	SSC CGL (Tier-I)	2019	22	$22 \times 25 = 550$
17.	SSC MTS	2019	39	$39 \times 25 = 975$
18.	SSC GD	2019	40	$40 \times 25 = 1000$
19.	SSC CHSL	2019	25	$25 \times 25 = 625$
20.	SSC CGL (Tier-II)	2017	7	$7 \times 100 = 700$
21.	SSC MTS	2017	17	$17 \times 25 = 425$
	Total		353	10,150



Note-Chapterwise compilation of total 10,150 questions of Quantitative Aptitude has been presented out of total 353 question papers of total 21 examinations conducted by Staff Selection Commission (SSC). Out of total 10150 questions asked from Quantitative Aptitude, total 5113 question of General Behavior have been removed and chapterwise compilation of 5037 questions of different types has been presented. In this book, every effort has been made by the Examination Special Committee to accommodate maximum variety of questions, so that examines can be made aware of the variety of questions asked by SSC.

Trend Analysis of Previous Year SSC Exams Papers Through Pie Chart and Bar Graph

SECTION-1



SECTION-2



(I) Problems based on Linear Equations

1. If $2x + 3y - 5z = 18$, $3x + 2y + z = 29$ and $x + y + 3z = 17$, then what is the value of $xy + yz + zx$?

- (a) 32 (b) 52
(c) 64 (d) 46

SSC CGL (Tier-II) 21-02-2018

Ans. (b) : Given,

$$2x + 3y - 5z = 18 \dots\dots (i)$$

$$3x + 2y + z = 29 \dots\dots (ii)$$

$$x + y + 3z = 17 \dots\dots (iii)$$

Multiplying by 5 in equation (ii) and adding it to equation (i),

$$2x + 3y - 5z = 18$$

$$15x + 10y + 5z = 145$$

$$17x + 13y = 163 \dots\dots (iv)$$

Again, on multiplying by 3 in equation (ii) and subtracting it to equation (iii) get-

$$8x + 5y = 70 \dots\dots (v)$$

By solving the equation (iv) and (v)

$$x = 5, y = 6$$

On putting the value of $x = 5$ and $y = 6$ in equation (ii),

$$15 + 12 + z = 29$$

$$\Rightarrow z = 2$$

$$\therefore xy + yz + zx = (5 \times 6) + (6 \times 2) + (2 \times 5)$$

$$= 30 + 12 + 10 = \boxed{52}$$

2. If $a - b = 3$ and $a^3 - b^3 = 999$, then find the value of $a^2 - b^2$.

- (a) 60 (b) 62
(c) 64 (d) 63

SSC CHSL 03/06/2022 (Shift- II)

Ans. (d) : $a - b = 3$

$$a^3 - b^3 = 999$$

$$(a - b) [(a - b)^2 + 3ab] = 999$$

$$3[9 + 3ab] = 999$$

$$3ab = 333 - 9$$

$$ab = 108$$

$$(a + b)^2 = (a - b)^2 + 4ab$$

$$(a + b)^2 = 9 + 4 \times 108$$

$$(a + b)^2 = 441$$

$$a + b = 21$$

$$a^2 - b^2 = (a - b)(a + b)$$

$$= 3 \times 21 = 63$$

3. If x, y, z are three integers such that $x + y = 8$, $y + z = 13$ and $z + x = 17$, then the value of $\frac{x^2}{yz}$ is:

- (a) 1 (b) $\frac{18}{11}$
(c) 0 (d) $\frac{7}{5}$

SSC CGL (Tier-I)-2019-03/03/2020 (Shift-I)

Ans. (b) : Given, $x + y = 8 \dots\dots (1)$

$$y + z = 13 \dots\dots (2)$$

$$z + x = 17 \dots\dots (3)$$

On adding the equation (i), (ii), and (iii)

$$2(x + y + z) = 38 \dots\dots (iv)$$

$$x + y + z = 19$$

$$\therefore x = 6, y = 2, z = 11$$

$$\therefore \frac{x^2}{yz} = \frac{36}{22} = \frac{18}{11}$$

Trick:

Put, $a = 6, y = 2, z = 11$

$$\therefore \frac{x^2}{y^2} = \frac{6^2}{2 \times 11} = \frac{18}{11}$$

4. If $3x + 6y + 9z = \frac{20}{3}$, $6x + 9y + 3z = \frac{17}{3}$ and $18x + 27y - z = \frac{113}{9}$, then what is the value of $75x + 113y$?

$$+ 113y ?$$

- (a) 163/3 (b) 143/6
(c) 218/9 (d) 311/3

SSC CGL (Tier-II) 9-3-2018

Ans. (a) :

$$3x + 6y + 9z = \frac{20}{3} \dots\dots (1)$$

$$6x + 9y + 3z = \frac{17}{3} \dots\dots (2)$$

$$18x + 27y - z = \frac{113}{9} \dots\dots (3)$$

On Multiplying by 3 in equation (iii) and adding it in equation (ii),

$$54x + 81y - 3z + 6x + 9y + 3z = \frac{113}{3} + \frac{17}{3}$$

$$60x + 90y = \frac{130}{3}$$

$$6x + 9y = \frac{13}{3} \dots\dots (4)$$

Multiplying by 3 in equation (2) and subtracting it from equation 1,

$$3x + 6y + 9z - 18x - 27y - 9z = \frac{20}{3} - 17$$

$$-15x - 21y = \frac{-31}{3}$$

$$15x + 21y = \frac{31}{3} \dots\dots (5)$$

Multiplying by 5 and 2 in equation (4) and equation (5) respectively then subtracting equation (5) from equation (4).

$$30x + 45y = \frac{65}{3}$$

$$30x + 42y = \frac{62}{3}$$

$$\frac{\quad}{3y = 1}$$

$$y = \frac{1}{3}$$

From equation (4),

$$6x + 3 = \frac{13}{3}$$

$$6x = \frac{13}{3} - 3$$

$$6x = \frac{4}{3}$$

$$x = \frac{2}{9}$$

$$\therefore 75x + 113y = 75 \times \frac{2}{9} + 113 \times \frac{1}{3}$$

$$= \frac{50}{3} + \frac{113}{3} = \frac{163}{3}$$

Trick:

$$(eq^n (i) + eq^n (ii)) \times \frac{1}{3} + eq^n (iii) \times 4$$

$$(9x + 15y + 12z) \times \frac{1}{3} + (72x + 108y - 4z)$$

$$= \frac{37}{9} + \frac{452}{9}$$

$$75x + 113y = \frac{489}{9} = \frac{163}{3}$$

5. If $3x + 4y - 2z + 9 = 17$, $7x + 2y + 11z + 8 = 23$ and $5x + 9y + 6z - 4 = 18$, then what is the value of $x + y + z - 34$?

(a) -28 (b) -14
(c) -31 (d) -45

SSC CGL (Tier-II) 20-02-2018

Ans. (c) : $3x + 4y - 2z + 9 = 17$ (i)
 $7x + 2y + 11z + 8 = 23$ (ii)
 $5x + 9y + 6z - 4 = 18$ (iii)

By adding the equation (i), (ii) and (iii)

$$15x + 15y + 15z = 45$$

$$x + y + z = 3$$

$$\therefore x + y + z - 34 = 3 - 34 = -31$$

6. If $x + 3y - \frac{2z}{4} = 6$, $x + \frac{2}{3}(2y + 3z) = 33$ and $\frac{1}{7}(x + y + z) + 2z = 9$, then what is the value of $46x + 131y$?

(a) 414 (b) 364
(c) 384 (d) 464

SSC CGL (Tier-II) 20-02-2018

Ans. (a) $x + 3y - \frac{2z}{4} = 6$
 $4x + 12y - 2z = 24$ -----(i)
 $x + \frac{2}{3}(2y + 3z) = 33$

$$3x + 4y + 6z = 99$$
 -----(ii)

$$\frac{1}{7}(x + y + z) + 2z = 9$$

$$x + y + z + 14z = 63$$

$$x + y + 15z = 63$$
 -----(iii)

From Equation (i) $\times \frac{21}{2}$ + Equation (ii) + Equation (iii),

$$42x + 126y - 21z + 3x + 4y + 6z + x + y + 15z = 252 + 99 + 63$$

$$46x + 131y = 414$$

7. If $3x + 4y - 11 = 18$ and $8x - 6y + 12 = 6$, then what is the value of $5x - 3y - 9$?

(a) 18 (b) -9
(c) -27 (d) -18

SSC CGL (Tier-II) 19-02-2018

Ans. (b) : $3x + 4y = 29$ (i)
 $8x - 6y = -6$ (ii)
 $4x - 3y = -3$ (ii)

On solving the equation (i) and (ii),
 $x = 3, y = 5$
 $\therefore 5x - 3y - 9 = 15 - 15 - 9 = -9$

8. If $a + b + c = 7/12$, $3a - 4b + 5c = 3/4$ and $7a - 11b - 13c = -7/12$, then what is the value of $a + c$?

(a) 1/2 (b) 5/12
(c) 3/4 (d) 1/4

SSC CGL (Tier-II) 19-02-2018

Ans. (b) : $a + b + c = \frac{7}{12}$ (1)
 $3a - 4b + 5c = \frac{3}{4}$ (2)
 $7a - 11b - 13c = -\frac{7}{12}$ (3)

Multiplying by 4 in equation (1) then adding it in equation (2)

$$4a + 4b + 4c + 3a - 4b + 5c = \frac{7}{3} + \frac{3}{4}$$

$$7a + 9c = \frac{37}{12}$$
(4)

On multiplying by 11 in equation (1) then adding it in equation (3)

$$11a + 11b + 11c + 7a - 11b - 13c = \frac{77}{12} - \frac{7}{12}$$

$$18a - 2c = \frac{35}{6}$$

$$9a - c = \frac{35}{12}$$
(5)

On multiplying by 9 in equation (5) then adding it in equation (4),

$$81a - 9c + 7a + 9c = \frac{315}{12} + \frac{37}{12}$$

$$88a = \frac{352}{12}$$

$$a = \frac{1}{3}$$

From equation (5),

$$3 - c = \frac{35}{12}$$

$$c = \frac{1}{12}$$

$$\text{thus } a + c = \frac{1}{3} + \frac{1}{12} = \frac{4+1}{12} = \frac{5}{12}$$

9. If $x - 4y = 0$ and $x + 2y = 24$, then what is the value of $(2x + 3y)/(2x - 3y)$?

- (a) $9/5$ (b) $11/5$
(c) $13/7$ (d) $9/7$

SSC CGL (Tier-II) 18-02-2018

Ans. (b):
Given,
 $x - 4y = 0$ (i)
 $x + 2y = 24$ (ii)
On putting the value $x = 4y$ in equation (ii).....
 $6y = 24$
 $y = 4$
 $\therefore x = 16$
As per question,
 $\frac{2x + 3y}{2x - 3y} = \frac{32 + 12}{32 - 12} = \frac{44}{20} = \frac{11}{5}$

10. If $3x + 5y + 7z = 49$ and $9x + 8y + 21z = 126$, then what is the value of y ?

- (a) 4 (b) 2
(c) 3 (d) 5

SSC CGL (Tier-II) 17-2-2018

Ans. (c) : $3x + 5y + 7z = 49$ (i)
 $9x + 8y + 21z = 126$ (ii)
On multiplying by 3 in equation (i),
 $9x + 15y + 21z = 147$ (iii)
From equation (iii) – Equation (ii)
 $7y = 21$
 $y = 3$

11. Cost of 8 pencils, 5 pens and 3 erasers is Rs. 111. Cost of 9 pencils, 6 pens and 5 erasers is Rs. 130. Cost of 16 pencils, 11 pens and 3 erasers is Rs. 221. What is the cost (in Rs.) of 39 pencils 26 pens and 13 erasers ?

- (a) 316 (b) 546
(c) 624 (d) 482

SSC CGL (Tier-II) 21-02-2018

Ans. (b) : Let, the price of 1 pencil, 1 pen and 1 eraser are x , y and z respectively.
As per question,
 $8x + 5y + 3z = 111$ (i)
 $9x + 6y + 5z = 130$ (ii)
 $16x + 11y + 3z = 221$ (iii)
By adding equations (i), (ii) and (iii)
 $33x + 22y + 11z = 462$ (iv)
On dividing by 11,
 $3x + 2y + z = 42$ (v)
From equation (v) $\times 13$
 $39x + 26y + 13z = 546$

12. A man buys 2 apples and 3 kiwi fruits for ₹37. If he buys 4 apples and 5 kiwi fruits for ₹67, then what will be the total cost of 1 apple and 1 kiwi fruit?

- (a) ₹20 (b) ₹18
(c) ₹15 (d) ₹28

SSC CHSL –17/03/2020 (Shift-II)

Ans. (c) : Let the cost price of an apple and a kiwi be x and y respectively.

As per question,

$$2x + 3y = 37 \text{(i)}$$

$$4x + 5y = 67 \text{(ii)}$$

By solving the equation (i) and (ii),

$$4x + 6y = 74 \text{ {multiplying by 2 in equation (i)}}$$

$$4x + 5y = 67$$

$$\begin{array}{r} - \quad - \quad - \\ \hline y = 7 \end{array}$$

From equation (i)–

$$2x + 3 \times 7 = 37$$

$$2x = 16$$

$$x = 8$$

Hence, the cost price of an apple and a kiwi = $8 + 7 = ₹15$

13. If $(1.25)(1 - 6.4 \times 10^{-5}) = 1.2496 + a$, then a equals :

- (a) 0.00032 (b) 0.0032
(c) 0.00016 (d) 0.0016

SSC CHSL (Tier-I) 11/07/2019 (Shift-II)

Ans. (a) : $(1.25)(1 - 6.4 \times 10^{-5}) = 1.2496 + a$
 $(1.25)\left(1 - 6.4 \times \frac{1}{10^5}\right) = 1.2496 + a$
 $(1.25)(1 - 0.000064) = 1.2496 + a$
 $1.25 \times 0.999936 = 1.2496 + a$
 $1.24992 - 1.2496 = a$
 $a = 0.00032$

14. If $u + v = 84$ and $u - v = 4$, then $u : v$ is equal to?

- (a) 11 : 10 (b) 10 : 11
(c) 10 : 9 (d) 9 : 10

SSC MTS 19/08/2019 (Shift-II)

Ans. (a) : Given,
 $u + v = 84$ (i)
 $u - v = 4$ (ii)
By solving the equation (i) and (ii)
 $2u = 88$
 $u = 44$
 $\therefore v = 40$
 $u : v = 44 : 40$
 $= 11 : 10$

15. The sum and difference of two numbers is 27 and 3 respectively. What is the ratio of two numbers?

- (a) 5 : 3 (b) 2 : 1
(c) 4 : 7 (d) 5 : 4

SSC MTS 16/08/2019 (Shift-III)

Ans. (d) : Let the numbers are x and y respectively.
As per question,
 $x + y = 27$ (i)
 $x - y = 3$ (ii)
By solving the equations (i) and (ii)
 $x = 15, y = 12$
Ratio of numbers $x : y = 15 : 12 = 5 : 4$

16. If the difference of two numbers is 7 and the difference of their squares is 203, then what is the smaller number?
 (a) 10 (b) 9
 (c) 12 (d) 11

SSC MTS 9-10-2017 (Shift-II)

Ans. (d) : Let the numbers are x and y

As per question,

$$x - y = 7 \text{ ----- (i)}$$

$$x^2 - y^2 = 203$$

$$(x+y)(x-y) = 203$$

$$7(x+y) = 203$$

$$x + y = 29 \text{(ii)}$$

By solving the equation (i) and (ii)

$$x = 18, y = 11$$

(II) Problems based on Algebraic Identities

17. If $2x - y = 2$ and $xy = \frac{3}{2}$, then what is the value of $x^3 - \frac{y^3}{8}$?

- (a) $\frac{9}{2}$ (b) $-\frac{5}{4}$
 (c) $\frac{5}{2}$ (d) $\frac{13}{4}$

SSC CGL (Tier-II) 29/01/2022

Ans : (d) Given, $2x - y = 2$ ------(i)

$$xy = \frac{3}{2}, x^3 - \frac{y^3}{8} = ?$$

On dividing by 2 and cubing both sides in eqⁿ, (i)

$$\left(x - \frac{y}{2}\right)^3 = (1)^3$$

$$x^3 - \frac{y^3}{8} - 3 \times \frac{xy}{2} \times 1 = 1$$

$$x^3 - \frac{y^3}{8} = \frac{13}{4}$$

18. If $x + \frac{16}{x} = 8$, then the value of $x^2 + \frac{32}{x^2}$ is:

- (a) 24 (b) 16
 (c) 20 (d) 18

SSC CGL (Tier-II)-2019-18/11/2020

Ans. (d) : $x + \frac{16}{x} = 8$ ------(Given)

$$\therefore x^2 - 8x + 16 = 0$$

$$(x - 4)^2 = 0$$

$$x = 4$$

$$\text{Hence, } x^2 + \frac{32}{x^2}$$

$$= 4^2 + \frac{32}{4^2} = 16 + \frac{32}{16}$$

$$= 16 + 2 = 18$$

Trick:

Put, $x = 4$

$$x + \frac{16}{x} = 8 \text{ (equation satisfies)}$$

$$\therefore x^2 + \frac{32}{x^2} = 16 + 2 = 18$$

19. If $(5\sqrt{5}x^3 - 81\sqrt{3}y^3) \div (\sqrt{5}x - 3\sqrt{3}y) = (Ax^2 + By^2 + Cxy)$, then the value of $(6A + B - \sqrt{15}C)$ is?
 (a) 10 (b) 15
 (c) 9 (d) 12

SSC CGL (TIER-I)-2018 - 04.06.2019 (Shift-I)

Ans. (d) :

$$\frac{5\sqrt{5}x^3 - 81\sqrt{3}y^3}{\sqrt{5}x - 3\sqrt{3}y} = Ax^2 + By^2 + Cxy$$

$$a^3 - b^3 = (a - b)(a^2 + b^2 + ab)$$

$$\frac{(\sqrt{5}x - 3\sqrt{3}y)(5x^2 + 27y^2 + 3\sqrt{15}xy)}{(\sqrt{5}x - 3\sqrt{3}y)} = Ax^2 + By^2 + Cxy$$

$$5x^2 + 27y^2 + 3\sqrt{15}xy = Ax^2 + By^2 + Cxy$$

By comparing the coefficient of x^2 , y^2 and xy , we get-

$$A = 5, B = 27, C = 3\sqrt{15}$$

$$6A + B - \sqrt{15}C = 30 + 27 - \sqrt{15} \times \sqrt{15} \times 3 = 57 - 45 = 12$$

20. If $x + 2y = 10$ and $2xy = 9$, then one of the value of $x - 2y$ is:

- (a) 8 (b) 6
 (c) 10 (d) 12

SSC CHSL 10/06/2022 (Shift- II)

Ans. (a) : Given that -

From formula -

$$(a + b)^2 - (a - b)^2 = 4ab$$

$$(10)^2 - (x - 2y)^2 = 4 \times 9$$

$$100 - 36 = (x - 2y)^2$$

$$64 = (x - 2y)^2$$

$$\therefore x - 2y = 8$$

Hence, option (a) is correct.

21. If $4x - 3y = 12$ and $xy = 5$, then find the value of $16x^2 + 9y^2$

- (a) 33 (b) 18
 (c) 3 (d) 44

SSC CGL (Tier-I) 21/04/2022 (Shift-III)

Ans : (c) $4x - 3y = 12$

On squaring both side

$$16x^2 + 9y^2 - 24xy = 144$$

$$16x^2 + 9y^2 = 144 - 120 \text{ (divide by 8 on both side)}$$

$$16x^2 + 9y^2 = 24$$

$$\frac{16x^2 + 9y^2}{8} = 3$$

22. If $\left(x + \frac{1}{x}\right) = \frac{11}{5}$, what is the value of

$$\left(x^3 + \frac{1}{x^3}\right)?$$

- (a) $4\frac{6}{125}$ (b) $5\frac{101}{125}$
 (c) $10\frac{81}{125}$ (d) $17\frac{31}{125}$

SSC CHSL 26/05/2022 (Shift- III)

Ans. (a) : Given,

$$x + \frac{1}{x} = \frac{11}{5}, \quad x^3 + \frac{1}{x^3} = ?$$

We know that :-

$$\text{If } x + \frac{1}{x} = a, \text{ then } x^3 + \frac{1}{x^3} = a^3 - 3a$$

$$\therefore a = 11/5$$

$$\therefore x^3 + \frac{1}{x^3} \Rightarrow \left(\frac{11}{5}\right)^3 - 3 \times \frac{11}{5}$$

$$\Rightarrow \frac{1331}{125} - \frac{33}{5} \Rightarrow \frac{1331 - 825}{125}$$

$$\Rightarrow \frac{506}{125} = 4\frac{6}{125}$$

23. If $x + \frac{1}{x} = -2\sqrt{3}$, what is the value of $x^5 + \frac{1}{x^5}$?

- (a) $-178\sqrt{3}$ (b) $-182\sqrt{3}$
 (c) $182\sqrt{3}$ (d) $-180\sqrt{3}$

SSC CHSL 24/05/2022 (Shift- III)

Ans. (a) : Given,

$$x + \frac{1}{x} = -2\sqrt{3} \text{ ----- (i)}$$

$$x^5 + \frac{1}{x^5} = ?$$

On cubing both sides of equation (i), we get-

$$x^3 + \frac{1}{x^3} + 3x \cdot \frac{1}{x} \left(x + \frac{1}{x}\right) = -8 \times 3\sqrt{3}$$

$$x^3 + \frac{1}{x^3} - 6\sqrt{3} = -24\sqrt{3}$$

$$\Rightarrow x^3 + \frac{1}{x^3} = -18\sqrt{3} \text{ (ii)}$$

Again on squaring both sides of equation (i), we get-

$$x^2 + \frac{1}{x^2} + 2 = 12$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 10 \text{ (iii)}$$

\therefore We know that,

$$x^5 + \frac{1}{x^5} = \left(x^3 + \frac{1}{x^3}\right) \left(x^2 + \frac{1}{x^2}\right) - \left(x + \frac{1}{x}\right)$$

From equation (i), (ii) & (iii), we get-

$$\begin{aligned} x^5 + \frac{1}{x^5} &= (-18\sqrt{3})(10) - (-2\sqrt{3}) \\ &= -180\sqrt{3} + 2\sqrt{3} \\ &= -178\sqrt{3} \\ \Rightarrow x^5 + \frac{1}{x^5} &= -178\sqrt{3} \end{aligned}$$

24. If $\left(a + \frac{1}{a} + 3\right) = 6$ where a is a non-zero real

number, then find the value of $a^2 + \frac{1}{a^2}$.

- (a) 3 (b) 47
 (c) 49 (d) 7

SSC CGL (Tier-I) 21/04/2022 (Shift-III)

$$\text{Ans : (d) } \left(a + \frac{1}{a} + 3\right) = 6$$

$$a + \frac{1}{a} = 3$$

On squaring both sides

$$a^2 + \frac{1}{a^2} + 2 \times a \times \frac{1}{a} = 3^2$$

$$a^2 + \frac{1}{a^2} = 9 - 2$$

$$a^2 + \frac{1}{a^2} = 7$$

25. If $a^2 + b^2 = 65$ and $ab = 8$, $a > b > 0$, then find the value of $a^2 - b^2$.

- (a) 72 (b) 63
 (c) 65 (d) 53

SSC CGL (Tier-I) 21/04/2022 (Shift-II)

Ans : (b) Given,

$$a^2 + b^2 = 65, ab = 8, a^2 - b^2 = ?$$

$$\begin{aligned} \text{Put, } a &= 8, b = 1 \\ a^2 - b^2 &= 8^2 - 1^2 \\ &= 64 - 1 = 63 \end{aligned}$$

26. If $x^4 + x^{-4} = 194$, $x > 0$ then the value of $(x-2)^2$ is?

- (a) 6 (b) 3
 (c) 2 (d) 1

SSC CGL (TIER-I)-2018 - 04.06.2019 (Shift-I)

$$\text{Ans. (b) } x^4 + \frac{1}{x^4} = 194$$

On adding 2 in both side

$$x^4 + \frac{1}{x^4} + 2 = 194 + 2$$

$$\left(x^2 + \frac{1}{x^2}\right)^2 = 196$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 14$$

On adding 2 in both sides

$$x^2 + \frac{1}{x^2} + 2 = 14 + 2$$

$$\left(x + \frac{1}{x}\right)^2 = 16$$

$$\Rightarrow x + \frac{1}{x} = 4$$

$$x^2 + 1 = 4x$$

$$x^2 - 4x + 1 = 0$$

On adding 3 in both sides

$$x^2 - 4x + 1 + 3 = 3$$

$$x^2 - 4x + 4 = 3$$

$$\boxed{(x-2)^2 = 3}$$

27. The expression $(a + b - c)^3 + (a - b + c)^3 - 8a^3$ is equal to:

- (a) $6a(a-b+c)(c-a-b)$ (b) $3a(a+b-c)(a-b+c)$
 (c) $6a(a+b-c)(a-b+c)$ (d) $3a(a-b+c)(c-a-b)$

SSC CGL (Tier-I)-2019 - 03/03/2020 (Shift-II)

Ans. (a) : $(a+b-c)^3 + (a-b+c)^3 - 8a^3$
 $= (a+b-c)^3 + (a-b+c)^3 + (-2a)^3 = 0$
 $\therefore A + B + C = a + b - c + a - b + c - 2a = 0$
 $\therefore A^3 + B^3 + C^3 = 3ABC$
 $\therefore (a+b-c)^3 + (a-b+c)^3 + (-2a)^3 = 3(a+b-c)(a-b+c)(-2a)$
 $= 6a(a-b+c)(c-a-b)$

Trick:

$$a = b + c = 1$$

Taken,

$$(a + b - c)^3 + (a - b + c)^3 - (2a)^3 = -6$$

From option (a)

$$6a(a - b + c)(c - a - b) = -6 \text{ equation satisfy}$$

28. If $x^4 - 79x^2 + 1 = 0$, then the value of $x + x^{-1}$ can be:

- (a) 9 (b) 5
(c) 7 (d) 8

SSC CGL (Tier-I) 21/04/2022 (Shift-II)

Ans : (a) $x^4 - 79x^2 + 1 = 0$
 On dividing by x^2 on both sides
 $x^2 - 79 + \frac{1}{x^2} = 0, x^2 + \frac{1}{x^2} = 79$
 Add 2 on both sides,
 $x^2 + \frac{1}{x^2} + 2 = 79 + 2$
 $\left(x + \frac{1}{x}\right)^2 = 81, x + \frac{1}{x} = 9$

29. If $a^3 + 3a^2 + 9a = 1$, then what is the value of $a^3 + (3/a)$?

- (a) 31 (b) 26
(c) 28 (d) 24

SSC CGL (Tier-II) 17-2-2018

Ans. (c) : $a^3 + 3a^2 + 9a = 1$
 $a^2 + 3a + 9 = \frac{1}{a}$
 Multiplying by $(a-3)$ in both side
 $(a-3)(a^2 + a \times 3 + 3^2) = \frac{1}{a} \times (a-3)$
 $a^3 - 3^3 = \frac{a-3}{a}$
 $a^3 + \frac{3}{a} = 1 + 27 = 28$

30. If $5x - \frac{1}{4x} = 6, x > 0$, then find the value of $25x^2 - \frac{1}{16x^2}$.

- (a) $6\sqrt{41}$ (b) 36
(c) $\sqrt{246}$ (d) $6\sqrt{31}$

SSC CGL (Tier-I) 21/04/2022 (Shift-I)

Ans : (a) Given-

$$5x - \frac{1}{4x} = 6 \text{ ----- (i)}$$

From formula $(a + b)^2 = (a - b)^2 + 4ab$

$$\therefore \left(5x + \frac{1}{4x}\right)^2 = \left(5x - \frac{1}{4x}\right)^2 + 4 \times 5x \times \frac{1}{4x}$$

$$\Rightarrow \left(5x + \frac{1}{4x}\right)^2 = \left(5x - \frac{1}{4x}\right)^2 + 5$$

$$\Rightarrow \left(5x + \frac{1}{4x}\right)^2 = (6)^2 + 5$$

$$\left(5x + \frac{1}{4x}\right) = \sqrt{41} \text{ ----- (ii)}$$

$$\therefore 25x^2 - \frac{1}{16x^2} = \left(5x - \frac{1}{4x}\right) \left(5x + \frac{1}{4x}\right)$$

$$= 6\sqrt{41} \quad \{\text{from eq}^n \text{ (i) \& (ii)}\}$$

31. If $x + y + z = 2, xy + yz + zx = -11$, and $xyz = -12$, then what is the value of $x^3 + y^3 + z^3$?

- (a) 36 (b) 38
(c) 40 (d) 42

SSC CGL (Tier-I) 13/04/2022 (Shift-III)

Ans : (b) Given,

$$x + y + z = 2, xy + yz + zx = -11, xyz = -12$$

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z) [(x + y + z)^2 - 3(xy + yz + zx)]$$

$$x^3 + y^3 + z^3 - 3 \times (-12) = 2[4 + 33]$$

$$x^3 + y^3 + z^3 = 74 - 36$$

$$x^3 + y^3 + z^3 = 38$$

32. If $2\sqrt{2}x^3 - 3\sqrt{3}y^3 = (\sqrt{2}x - \sqrt{3}y)(Ax^2 - Bxy + Cy^2)$, then the value of $(A^2 + B^2 + C^2)$ is:

- (a) 16 (b) 11
(c) 19 (d) 18

SSC CGL (Tier-I) 13/04/2022 (Shift-II)

Ans : (c)

$$2\sqrt{2}x^3 - 3\sqrt{3}y^3 = (\sqrt{2}x - \sqrt{3}y)(Ax^2 - Bxy + Cy^2)$$

$$(\sqrt{2}x - \sqrt{3}y)(2x^2 + \sqrt{6}xy + 3y^2) = (\sqrt{2}x - \sqrt{3}y)(Ax^2 - Bxy + Cy^2)$$

$$Ax^2 - Bxy + Cy^2 = (2x^2 + \sqrt{6}xy + 3y^2)$$

$$A = 2, B = -\sqrt{6}, C = 3$$

$$A^2 + B^2 + C^2 = 2^2 + 6 + 3^2$$

$$= 4 + 6 + 9$$

$$= 19$$

33. If $a + b + c = 11$ and $ab + bc + ca = 28$, then find the value of $a^3 + b^3 + c^3 - 3abc$.
- (a) 1639 (b) 407
(c) 2255 (d) 1093

SSC CGL (Tier-I) 19/04/2022 (Shift-III)

Ans. (b) $a + b + c = 11$, $ab + bc + ca = 28$
 $a^3 + b^3 + c^3 - 3abc = (a + b + c) [(a + b + c)^2 - 3(ab + bc + ca)]$
 $a^3 + b^3 + c^3 - 3abc = 11(121 - 84)$
 $a^3 + b^3 + c^3 - 3abc = 11 \times 37$
 $a^3 + b^3 + c^3 - 3abc = 407$

34. If $a^2 + b^2 + 49c^2 + 18 = 2(b + 28c - a)$, then the value of $(2a - b + 7c)$ is:
- (a) 5 (b) -3
(c) -4 (d) 1

SSC CGL (Tier-I) 19/04/2022 (Shift-II)

Ans. (d) Given,
 $a^2 + b^2 + 49c^2 + 18 = 2(b + 28c - a)$
 $a^2 + b^2 + 49c^2 + 18 = 2b + 56c - 2a$
 $(a^2 + 2a + 1) + (b^2 - 2b + 1) + (49c^2 - 56c + 16) = 0$
 $(a + 1)^2 + (b - 1)^2 + (7c - 4)^2 = 0$
 $a + 1 = 0, b - 1 = 0, 7c - 4 = 0$
 $a = -1, b = 1, c = \frac{4}{7}$
 $\therefore (2a - b + 7c) = 2 \times (-1) - (1) + 7 \times \frac{4}{7} = 1$

35. If $x + y + z = 7$, $xy + yz + zx = 8$, then what is the value of $x^3 + y^3 + z^3 - 3xyz$?
- (a) 200 (b) 150
(c) 125 (d) 175

SSC CGL (Tier-I) 19/04/2022 (Shift-I)

Ans. (d) Given, $x + y + z = 7$, $xy + yz + zx = 8$
 Now,
 $x^3 + y^3 + z^3 - 3xyz = (x + y + z) [(x + y + z)^2 - 3(xy + yz + zx)]$
 $= 7 [49 - 3 \times 8]$
 $= 7 \times 25$
 $= 175$

36. If $\sqrt{x} - \frac{1}{\sqrt{x}} = \sqrt{3}$, then what is the value of

$$x^4 + \frac{1}{x^4} ?$$

- (a) 531 (b) 7
(c) 623 (d) 527

SSC CGL (Tier-I) 18/04/2022 (Shift-III)

Ans. (d) From question,

$$\sqrt{x} - \frac{1}{\sqrt{x}} = \sqrt{3}$$

On squaring both sides,

$$\left(\sqrt{x} - \frac{1}{\sqrt{x}}\right)^2 = (\sqrt{3})^2$$

$$\Rightarrow x + \frac{1}{x} - 2 \times \sqrt{x} \times \frac{1}{\sqrt{x}} = 3$$

$$\Rightarrow x + \frac{1}{x} = 3 + 2$$

$$\Rightarrow x + \frac{1}{x} = 5$$

Again, on squaring both sides

$$\left(x + \frac{1}{x}\right)^2 = 5^2$$

$$x^2 + \frac{1}{x^2} + 2 = 25$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 23$$

Again, squaring on both sides

$$\left(x^2 + \frac{1}{x^2}\right)^2 = (23)^2$$

$$\Rightarrow x^4 + \frac{1}{x^4} + 2 = 529$$

$$\therefore x^4 + \frac{1}{x^4} = 527$$

37. If $2\sqrt{2}x^3 - 3\sqrt{3}y^3 = (\sqrt{2}x - \sqrt{3}y)(Ax^2 - Bxy + Cy^2)$, then the value of $\sqrt{(A^2 + B^2 + C^2)}$ is:

- (a) $\sqrt{19}$ (b) $\sqrt{11}$
(c) $\sqrt{17}$ (d) $\sqrt{21}$

SSC CGL (Tier-I) 18/04/2022 (Shift-II)

Ans. (a) $2\sqrt{2}x^3 - 3\sqrt{3}y^3 = (\sqrt{2}x - \sqrt{3}y)(Ax^2 - Bxy + Cy^2)$,

$$(\sqrt{2}x - \sqrt{3}y)(2x^2 + \sqrt{6}xy + 3y^2) = (\sqrt{2}x - \sqrt{3}y)(Ax^2 - Bxy + Cy^2)$$

$$(2x^2 + \sqrt{6}xy + 3y^2) = (Ax^2 - Bxy + Cy^2)$$

On comparing,

$$A = 2, B = -\sqrt{6}, C = 3$$

$$\sqrt{(A^2 + B^2 + C^2)} = \sqrt{2^2 + (-\sqrt{6})^2 + (3)^2} = \sqrt{4 + 6 + 9} = \sqrt{19}$$

38. If $a^2 + b^2 + 49c^2 + 18 = 2(b - 28c - a)$, then the value of $(a - b - 7c)$ is:

- (a) 4 (b) 3
(c) 2 (d) 1

SSC CGL (Tier-I) 18/04/2022 (Shift-II)

Ans. (c) $a^2 + b^2 + 49c^2 + 18 = 2(b - 28c - a)$

$$(a + 1)^2 + (b - 1)^2 + (7c + 4)^2 = 0$$

$$a = -1, b = 1, c = -4/7$$

$$\text{Now, } (a - b - 7c) = (-1 - 1 + 7 \times \frac{4}{7}) = (-2 + 4) = 2$$

39. If $\left(x^2 + \frac{1}{x^2}\right) = 23$, $x > 0$ What is the value of

$$\left(x^3 + \frac{1}{x^3}\right) = ?$$

- (a) 140 (b) 110
(c) -110 (d) -140

SSC CGL (Tier-I) 18/04/2022 (Shift-I)

Ans. (b) $\left(x^2 + \frac{1}{x^2}\right) = 23$

On adding 2 both sides,

$$x^2 + \frac{1}{x^2} + 2 = 23 + 2$$

$$= \left(x + \frac{1}{x}\right)^2 = 5^2$$

$$x + \frac{1}{x} = 5$$

On cubing both sides

$$x^3 + \frac{1}{x^3} + 3 \times 5 = 125$$

$$\therefore x^3 + \frac{1}{x^3} = (5)^3 - 3 \times 5 = 125 - 15 = 110$$

40. If $x + y + z = 18$, $xyz = 81$ and $xy + yz + zx = 90$, then the value of $x^3 + y^3 + z^3 + xyz$ is:

- (a) 1321 (b) 1296
(c) 1225 (d) 1250

SSC CGL (Tier-I) 13/04/2022 (Shift-I)

Ans. (b) Given,

$$x + y + z = 18, xyz = 81, xy + yz + zx = 90$$

From the formula,

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z) [(x + y + z)^2 - 3(xy + yz + zx)]$$

$$x^3 + y^3 + z^3 - 3xyz = 18 [(18)^2 - 3(90)]$$

$$x^3 + y^3 + z^3 + xyz = 18(324 - 270) + 4xyz$$

$$= 18 \times 54 + 4 \times 81$$

$$= 972 + 324$$

$$x^3 + y^3 + z^3 + xyz = 1296$$

41. If $a^2 + b^2 + c^2 = 6.25$ and $(ab + bc + ca) = 0.52$, what is the value of $(a+b+c)$, if $(a + b + c) < 0$?

- (a) ± 2.7 (b) -2.7
(c) -2.8 (d) ± 2.8

SSC CGL (Tier-I) 11/04/2022 (Shift-III)

Ans. (b) Given,

$$a^2 + b^2 + c^2 = 6.25$$

$$ab + bc + ca = 0.52$$

$$\text{Let } a + b + c = y$$

On squaring both sides,

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = y^2$$

$$6.25 + 2 \times (0.52) = y^2$$

$$6.25 + 1.04 = y^2$$

$$y^2 = 7.29$$

$$y = \pm \sqrt{7.29}$$

$$\text{If } a + b + c < 0$$

$$\text{Then, } a + b + c = -2.7$$

42. If $x + y + 3 = 0$, then find the value of $x^3 + y^3 - 9xy + 9$.

- (a) -18 (b) -36
(c) 18 (d) 36

SSC CGL (Tier-I) 11/04/2022 (Shift-I)

Ans. (a) $x + y + 3 = 0$

$$x + y = -3$$

On cubing both sides

$$(x + y)^3 = -3^3$$

$$x^3 + y^3 + 3xy(x + y) = -27$$

$$x^3 + y^3 + 3xy(-3) = -27$$

$$x^3 + y^3 - 9xy = -27$$

$$x^3 + y^3 - 9xy + 9 = -27 + 9 = -18$$

43. If $(4x + 2y)^3 + (4x - 2y)^3 = 16(Ax^3 + Bxy^2)$, then what is the value of $\frac{1}{2}(\sqrt{A^2 + B^2})$?

- (a) 8 (b) 3
(c) 5 (d) 7

SSC CGL (Tier-I) 11/04/2022 (Shift-II)

Ans. (c) $(4x + 2y)^3 + (4x - 2y)^3 = 16(Ax^3 + Bxy^2)$
 $64x^3 + 8y^3 + 24xy(4x + 2y) + 64x^3 - 8y^3 - 24xy(4x - 2y) = 16(Ax^3 + Bxy^2)$

$$128x^3 + 96xy^2 = 16(Ax^3 + Bxy^2)$$

$$16(8x^3 + 6xy^2) = 16(Ax^3 + Bxy^2)$$

On comparing both sides

$$A = 8 \quad B = 6$$

Then,

$$\frac{1}{2}(\sqrt{A^2 + B^2}) = \frac{1}{2}(\sqrt{8^2 + 6^2}) = 5$$

44. If $x = 4 + \sqrt{15}$, What is the value of

$$\left(x^2 + \frac{1}{x^2}\right)?$$

- (a) 48 (b) 54
(c) 72 (d) 62

SSC CGL (Tier-I) 11/04/2022 (Shift-III)

Ans. (d) Given,

$$x = 4 + \sqrt{15}$$

$$\frac{1}{x} = \frac{1}{4 + \sqrt{15}} = \frac{1}{4 + \sqrt{15}} \times \frac{4 - \sqrt{15}}{4 - \sqrt{15}} = 4 - \sqrt{15}$$

$$x + \frac{1}{x} = 4 + \sqrt{15} + 4 - \sqrt{15}$$

$$x + \frac{1}{x} = 8$$

On squaring both sides

$$x^2 + \frac{1}{x^2} + 2 \times x \times \frac{1}{x} = 64$$

$$x^2 + \frac{1}{x^2} = 62$$

45. If $x + \frac{1}{x} = 3$, $x \neq 0$, then the value of $x^7 + \frac{1}{x^7}$ is:

- (a) 749 (b) 843
(c) 746 (d) 849

SSC CGL (Tier-II) 03/02/2022

Ans : (b) $x + \frac{1}{x} = 3$

$$x^3 + \frac{1}{x^3} = 3^3 - 3 \times 3 = 18$$

Again,

$$x + \frac{1}{x} = 3$$

On squaring both sides

$$x^2 + \frac{1}{x^2} = 7$$

On squaring both sides

$$x^4 + \frac{1}{x^4} = 7^2 - 2 = 47$$

$$x^7 + \frac{1}{x^7} = \left(x^3 + \frac{1}{x^3}\right)\left(x^4 + \frac{1}{x^4}\right) - \left(x + \frac{1}{x}\right)$$

$$= (18)(47) - 3$$

$$= 846 - 3 = 843$$

46. If $x^2 - 3x + 1 = 0$, then the value of

$$\frac{\left(x^4 + \frac{1}{x^2}\right)}{(x^2 + 5x + 1)} \text{ is:}$$

- (a) $\frac{9}{4}$ (b) $\frac{27}{8}$
(c) $\frac{5}{2}$ (d) 2

SSC CGL (Tier-II) 03/02/2022

Ans : (a) $x^2 - 3x + 1 = 0$

$$x\left(x - 3 + \frac{1}{x}\right) = 0$$

$$x + \frac{1}{x} = 3$$

On cubing both sides,

$$x^3 + \frac{1}{x^3} + 3 \times 3 = 27$$

$$x^3 + \frac{1}{x^3} = 3^3 - 9 = 18$$

$$\frac{x^4 + \frac{1}{x^2}}{x^2 + 5x + 1}$$

$$x\left(x^3 + \frac{1}{x^3}\right)$$

$$x\left[\left\{x + \frac{1}{x}\right\} + 5\right]$$

$$\frac{18}{3+5} = \frac{9}{4}$$

47. If $a + b = 8$, $ab = 10$, then the value of $a^3 + b^3$ is:

- (a) 312 (b) 215
(c) 272 (d) 111

SSC CGL (Tier-II) 29/01/2022

Ans : (c) Given

$$a + b = 8$$

$$ab = 10$$

$$a^3 + b^3 = ?$$

we know that,

$$a^3 + b^3 = (a + b)[(a + b)^2 - 3ab]$$

$$= (8)[(8)^2 - 3 \times 10]$$

$$= 8[64 - 30]$$

$$= 8 \times 34 = 272$$

48. If $a + b + c = 1$, $ab + bc + ca = -22$ and $abc = -40$, then what is the value of $a^3 + b^3 + c^3$?

- (a) 67 (b) -53
(c) -51 (d) 27

SSC CGL (Tier-II) 29/01/2022

Ans : (b) Given,

$$a + b + c = 1$$

$$ab + bc + ca = -22$$

$$abc = -40$$

$$a^3 + b^3 + c^3 = ?$$

$$\therefore a^3 + b^3 + c^3 - 3abc = (a + b + c)[(a + b + c)^2 - 3(ab + bc + ca)]$$

$$a^3 + b^3 + c^3 + 120 = (1)[(1)^2 + 3(22)]$$

$$= 1 \times 67$$

$$\Rightarrow a^3 + b^3 + c^3 = 67 - 120$$

$$\Rightarrow a^3 + b^3 + c^3 = -53$$

49. If $27x^3 - 64y^3 = (Ax + By)(Cx^2 + Dy^2 - Exy)$, then value of $(A - B + C - D + E)$ will be:

- (a) -12 (b) 18
(c) 15 (d) -20

SSC CHSL 09/08/2021 (Shift-I)

Ans. (a) : $27x^3 - 64y^3 = (Ax + By)(Cx^2 + Dy^2 - Exy)$ --- (Given)

$$(3x - 4y)(9x^2 + 16y^2 + 12xy) = (Ax + By)(Cx^2 + Dy^2 - Exy)$$

On comparing both sides, we get,

$$A = 3, B = -4, C = 9, D = 16, E = -12$$

Hence

$$A - B + C - D + E = 3 + 4 + 9 - 16 + (-12)$$

$$= 16 - 16 - 12$$

$$= -12$$

50. If $(3x + 2y)^3 + (3x - 2y)^3 = 3kx(3x^2 + 4y^2)$, then the value of k will be:

- (a) 18 (b) 9
(c) 3 (d) 6

SSC CHSL 09/08/2021 (Shift-I)

Ans. (d) : $(3x + 2y)^3 + (3x - 2y)^3 = 3kx(3x^2 + 4y^2)$

$$\therefore a^3 + b^3 = (a+b)(a^2 - ab + b^2)$$

So,

$$(3x + 2y + 3x - 2y)[(3x + 2y)^2 + (3x - 2y)^2 - (3x + 2y)(3x - 2y)]$$

$$= 3kxy(3x^2 + 4y^2)$$

$$6x[(9x^2 + 4y^2) \times 2 - (9x^2 - 4y^2)] = 3kx(3x^2 + 4y^2)$$

$$6x[18x^2 + 8y^2 - 9x^2 + 4y^2] = 3kx(3x^2 + 4y^2)$$

$$6x[9x^2 + 12y^2] = 3kx(3x^2 + 4y^2)$$

$$3 \times 6x(3x^2 + 4y^2) = 3kx(3x^2 + 4y^2)$$

$$6x = kx$$

$$k = 6$$

51. If $x + 2y = 19$ and $x^3 + 8y^3 = 361$, then xy is equal to:

- (a) 57 (b) 56
(c) 55 (d) 58

SSC CHSL 09/08/2021 (Shift-I)

Ans. (a) : $x + 2y = 19$... (i)
 $x^3 + 8y^3 = 361$... (ii)

On cubing both sides of equation (i),

$$x^3 + 8y^3 + 6xy(x + 2y) = 6859$$

By equation (i) and (ii),

$$361 + 6xy(19) = 6859$$

$$6xy \times 19 = 6859 - 361 = 6498$$

$$xy = \frac{6498}{114} = 57$$

52. If $x^2 + 4y^2 + 3z^2 + \frac{19}{4} = 2\sqrt{3}(x+y+z)$, then the value of $(x - 4y + 3z)$ is:

- (a) $\frac{\sqrt{3}}{3}$ (b) $2\sqrt{3}$
 (c) $\sqrt{3}$ (d) $\frac{\sqrt{3}}{2}$

SSC CHSL 05/08/2021 (Shift-I)

Ans. (c) : Given,

$$x^2 + 4y^2 + 3z^2 + \frac{19}{4} = 2\sqrt{3}(x+y+z)$$

$$x^2 - 2\sqrt{3}x + 3 + 4y^2 - 2\sqrt{3}y + \frac{3}{4} + 3z^2 - 2\sqrt{3}z + 1 = 0$$

$$(x - \sqrt{3})^2 + \left(2y - \frac{\sqrt{3}}{2}\right)^2 + (\sqrt{3}z - 1)^2 = 0$$

$$x = \sqrt{3}, \quad 2y = \frac{\sqrt{3}}{2}, \quad \sqrt{3}z = 1, \Rightarrow y = \frac{\sqrt{3}}{4}, z = \frac{1}{\sqrt{3}}$$

Hence,

$$\begin{aligned} x - 4y + 3z &= \sqrt{3} - 4 \times \frac{\sqrt{3}}{4} + 3 \times \frac{1}{\sqrt{3}} \\ &= \sqrt{3} - \sqrt{3} + \sqrt{3} \\ &= \sqrt{3} \end{aligned}$$

53. If $x + y + z = 13$, $x^2 + y^2 + z^2 = 91$ and $xz = y^2$, then the difference between z and x is:

- (a) 3 (b) 8
 (c) 5 (d) 9

SSC CHSL 05/08/2021 (Shift-I)

Ans. (b) : Given,

$$x + y + z = 13 \quad x^2 + y^2 + z^2 = 91 \text{ and } xz = y^2$$

$$x + y + z = 13$$

On squaring both sides,

$$x^2 + y^2 + z^2 + 2(xy + yz + zx) = 169$$

$$91 + 2(xy + yz + y^2) = 169$$

$$2(x+y+z)y = 169 - 91$$

$$y = \frac{39}{13} = 3$$

$$x + z = 13 - 3$$

$$x + z = 10$$

$$xz = 9$$

$$z = \frac{9}{x}$$

$$x + \frac{9}{x} = 10$$

$$x(10-x) = 9$$

$$x = 1$$

Hence, $z - x = 9 - 1 = 8$

54. If $x + y = 5$ and $\frac{1}{x} + \frac{1}{y} = \frac{20}{9}$, then the value of $(x^3 + y^3)$ will be:

- (a) $\frac{635}{8}$ (b) $\frac{365}{4}$
 (c) $\frac{205}{4}$ (d) $\frac{635}{4}$

SSC CHSL 15/04/2021 (Shift-I)

Ans. (b) : $x + y = 5$ (i)

$$\frac{1}{x} + \frac{1}{y} = \frac{20}{9}$$

$$\frac{x+y}{xy} = \frac{20}{9}$$

$$\frac{5}{xy} = \frac{20}{9} \quad [\text{On putting the value of } x + y = 5]$$

$$xy = \frac{9}{4}$$

$$(x+y)^3 = x^3 + y^3 + 3xy(x+y)$$

$$(5)^3 = x^3 + y^3 + 3 \times \frac{9}{4} \times 5$$

$$125 - \frac{135}{4} = x^3 + y^3$$

$$x^3 + y^3 = \frac{500 - 135}{4} = \frac{365}{4}$$

55. If $x + y + z = 5$, $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 0$, $xyz = 12$ and $x^3 + y^3 + z^3 = 151$, then the value of $(x^2 + y^2 + z^2)$ is:

- (a) 23 (b) 24
 (c) 21 (d) 22

SSC CHSL 15/04/2021 (Shift-I)

Ans. (a) : Given $x+y+z = 5$, $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 0$, $xyz = 12$

$$x^3 + y^3 + z^3 = 151$$

$$\therefore \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 0$$

$$yz + zx + xy = 0$$

$$x^3 + y^3 + z^3 - 3xyz = (x+y+z)(x^2 + y^2 + z^2 - xy - yz - zx)$$

$$151 - 3 \times 12 = 5[x^2 + y^2 + z^2 - (xy + yz + zx)]$$

$$115 = 5(x^2 + y^2 + z^2 - 0)$$

$$x^2 + y^2 + z^2 = \frac{115}{5} = 23$$

56. If $49a^2 + 25b^2 = 30$ and $ab = 1$, $a, b > 0$, then the value of $(7a + 5b)$ is:

- (a) 14 (b) 10
 (c) 8 (d) 12

SSC CHSL 15/04/2021 (Shift-I)

Ans. (b) : $49a^2 + 25b^2 = 30$

On adding $70ab$ both sides,

$$49a^2 + 25b^2 + 70ab = 30 + 70ab$$

$$(7a)^2 + (5b)^2 + 2 \times 7 \times 5ab = 30 + 70ab \quad (\because ab = 1)$$

$$(7a + 5b)^2 = 30 + 70$$

$$7a + 5b = \sqrt{100} = 10$$

57. If $x^4 + \frac{1}{x^4} = 727, x > 1$, then what is the value of

$$\left(x - \frac{1}{x}\right)^2 ?$$

- (a) 6 (b) 5
(c) -5 (d) -6

SSC CGL-(Tier-I) 13/08/2021 (Shift III)

Ans. (b) : Given that- $x^4 + \frac{1}{x^4} = 727, x > 1$

Adding 2 on both sides, $x^4 + \frac{1}{x^4} + 2 = 727 + 2$

$$\Rightarrow \left(x^2 + \frac{1}{x^2}\right)^2 = (27)^2$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 27$$

Subtracting 2 from both sides,

$$x^2 + \frac{1}{x^2} - 2 = 27 - 2$$

$$\Rightarrow x^2 + \frac{1}{x^2} - 2 \times x \times \frac{1}{x} = 25$$

$$\Rightarrow \left(x - \frac{1}{x}\right)^2 = 5^2$$

Taking square root both sides,

$$\boxed{x - \frac{1}{x} = 5}$$

Hence, option (b) is correct.

58. If $x - \frac{1}{x} = 1$, then what is the value of $x^8 + \frac{1}{x^8}$?

- (a) 119 (b) -1
(c) 3 (d) 47

SSC CGL-(Tier-I) 13/08/2021 (Shift III)

Ans. (d) : From question,

$$x - \frac{1}{x} = 1$$

On squaring both sides,

$$\left(x - \frac{1}{x}\right)^2 = 1^2$$

$$\Rightarrow x^2 + \frac{1}{x^2} - 2 = 1$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 3$$

Again on squaring both sides,

$$\left(x^2 + \frac{1}{x^2}\right)^2 = 3^2$$

$$\Rightarrow x^4 + \frac{1}{x^4} + 2 = 9, \Rightarrow x^4 + \frac{1}{x^4} = 7$$

Again on squaring both sides,

$$\left(x^4 + \frac{1}{x^4}\right)^2 = 7^2$$

$$\Rightarrow x^8 + \frac{1}{x^8} + 2 = 49$$

$$\therefore \boxed{x^8 + \frac{1}{x^8} = 47}$$

59. If $2x^2 - 7x + 5 = 0$, then what is the value of

$$x^3 + \frac{125}{8x^3} ?$$

- (a) $12\frac{5}{8}$ (b) $16\frac{5}{8}$
(c) $10\frac{5}{8}$ (d) $18\frac{5}{8}$

SSC CGL-(Tier-I) 2308/2021 (Shift I)

Ans. (b) : $2x^2 - 7x + 5 = 0$

On dividing by $2x$, on both sides

$$x - \frac{7}{2} + \frac{5}{2x} = 0$$

$$x + \frac{5}{2x} = \frac{7}{2}$$

On cubing both sides,

$$x^3 + \frac{125}{8x^3} + 3 \times x \times \frac{5}{2x} \left(x + \frac{5}{2x}\right) = \frac{343}{8}$$

$$x^3 + \frac{125}{8x^3} + \frac{15}{2} \times \frac{7}{2} = \frac{343}{8}$$

$$x^3 + \frac{125}{8x^3} = \frac{343}{8} - \frac{105}{4} = \frac{133}{8}$$

$$= 16\frac{5}{8}$$

60. If $2x + 3y + 1 = 0$, then what is the value of $(8x^3 + 8 + 27y^3 - 18xy)$?

- (a) -7 (b) 7
(c) -9 (d) 9

SSC CGL-(Tier-I) 24/08/2021 (Shift I)

Ans. (b) : If $a + b + c = 0$

$$\therefore a^3 + b^3 + c^3 - 3abc = 0$$

$$\therefore 2x + 3y + 1 = 0$$

$$\therefore 8x^3 + 27y^3 + (1)^3 - 3 \times 2x \times 3y \times 1 = 0$$

$$8x^3 + 27y^3 + 1 - 18xy = 0 \text{ (adding +7 to Both sides)}$$

$$8x^3 + 27y^3 + 8 - 18xy = 7$$

61. If $a^4 + b^4 + a^2b^2 = 273$ and $a^2 + b^2 - ab = 21$,

then one of the values of $\left(\frac{1}{a} + \frac{1}{b}\right)$ is :

- (a) $-\frac{9}{4}$ (b) $-\frac{3}{4}$
(c) $\frac{9}{8}$ (d) $\frac{3}{2}$

SSC CGL-(Tier-I) 24/08/2021 (Shift I)

Ans. (b) : $\therefore (a^2 + b^2 + ab)(a^2 + b^2 - ab) = a^4 + b^4 + a^2b^2$

$$\therefore a^2 + b^2 + ab = \frac{273}{21} = 13 \text{(1)}$$

$$\text{Given } a^2 + b^2 - ab = 21 \text{(2)}$$

From eqⁿ (1) - (2),
 $2ab = -8$
 From eqⁿ (1) + (2),
 $2(a^2 + b^2) = 34$
 $a^2 + b^2 = 17$
 $\therefore (a+b)^2 = 17 - 8 = 9$
 $a + b = 3$
 $\therefore \frac{1}{a} + \frac{1}{b} = \frac{b+a}{ab} = \frac{3}{-4} = -\frac{3}{4}$

62. If $(54\sqrt{2}x^3 + 24\sqrt{3}y^3) \div (\sqrt{18}x + \sqrt{12}y) = Ax^2 + By^2 + Cxy$, then what is the value of $A^2 - (B^2 + C^2)$?

- (a) 12 (b) -36
 (c) -24 (d) 24

SSC CGL-(Tier-I) 17/08/2021 (Shift I)

Ans. (b) :

$$\frac{[(3\sqrt{2}x)^3 + (2\sqrt{3}y)^3] \div (3\sqrt{2}x + 2\sqrt{3}y)}{= Ax^2 + By^2 + Cxy}$$

$$\therefore a^3 + b^3 = (a+b)(a^2 + b^2 - ab)$$

$$\frac{(3\sqrt{2}x + 2\sqrt{3}y)(18x^2 + 12y^2 - 6\sqrt{6}xy)}{(3\sqrt{2}x + 2\sqrt{3}y)} = Ax^2 + By^2 + Cxy$$

On comparing:

$$Ax^2 + By^2 + Cxy = 18x^2 + 12y^2 - 6\sqrt{6}xy$$

A = 18, B = 12, C = $-6\sqrt{6}$
 $\therefore A^2 - (B^2 + C^2) = 18^2 - (144 + 216)$
 $= 324 - 360$
 $= -36$

63. If $x + y + z = 7$, $x^2 + y^2 + z^2 = 85$ and $x^3 + y^3 + z^3 = 913$, then the value of $\sqrt[3]{xyz}$ is:

- (a) 1 (b) 2
 (c) 4 (d) 8

SSC CGL-(Tier-I) 17/08/2021 (Shift I)

Ans. (c) : Given,

$$x + y + z = 7, x^2 + y^2 + z^2 = 85, x^3 + y^3 + z^3 = 913$$

$$\therefore x^3 + y^3 + z^3 - 3xyz = (x+y+z)[x^2 + y^2 + z^2 - xy - yz - zx]$$

$$\therefore (x+y+z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)$$

$$\therefore x^2 + y^2 + z^2 + 2(xy + yz + zx) = 49$$

$$2(xy + yz + zx) = -36$$

$$(xy + yz + zx) = -18$$

$$x^3 + y^3 + z^3 - 3xyz = 7 \times [85 + 18]$$

$$3xyz = 913 - 721 = 192$$

$$xyz = 64$$

$$\sqrt[3]{xyz} = \sqrt[3]{64} = 4$$

64. If $x^4 + y^4 + x^2y^2 = 21$ and $x^2 + y^2 - xy = 7$, then what is the value of $\frac{x}{y} + \frac{y}{x}$?

- (a) $\frac{5}{4}$ (b) $\frac{3}{4}$

- (c) $-\frac{3}{2}$ (d) $-\frac{5}{2}$

SSC CGL-(Tier-I) 18/08/2021 (Shift I)

Ans. (d) : $x^4 + y^4 + x^2y^2 = 21$
 given $x^2 + y^2 - xy = 7$ (i)
 $(x^2 + y^2 + xy)(x^2 + y^2 - xy) = x^4 + y^4 + x^2y^2$
 $(x^2 + y^2 + xy) \times 7 = 21$
 $x^2 + y^2 + xy = 3$(ii)
 Equation (i) + Equation (ii),
 $x^2 + y^2 + xy = 3$
 $\frac{x^2 + y^2 - xy = 7}{x^2 + y^2 + xy = 3}$
 $2(x^2 + y^2) = 10$
 $x^2 + y^2 = 5$
 $x^2 + y^2 + xy = 3$
 $xy = -2$
 $\frac{x}{y} + \frac{y}{x} = \frac{x^2 + y^2}{xy} = \frac{-5}{2}$

65. If $x - y = 11$ and $\frac{1}{x} - \frac{1}{y} = \frac{11}{24}$, then what is the value of $x^3 - y^3 + x^2y^2$?

- (a) 1331 (b) 1105
 (c) 1307 (d) 1115

SSC CGL-(Tier-I) 13/08/2021 (Shift II)

Ans. (d) Given,

$$x - y = 11 \text{ and } \frac{1}{x} - \frac{1}{y} = \frac{11}{24}, \frac{y-x}{xy} = \frac{11}{24}$$

$$xy = -1 \times 24$$

$$xy = -24$$

$$x^3 - y^3 + x^2y^2 = ?$$

$$= (x-y)[(x-y)^2 + 3xy] + (-24)^2$$

$$= 11 \times [121 - 72] + 576$$

$$= 11 \times 49 + 576$$

$$= 539 + 576 = 1115$$

66. If $(16\sqrt{2}x^3 + 81\sqrt{3}y^3) \div (2\sqrt{2}x + 3\sqrt{3}y) = Ax^2 + By^2 + Cxy$, then find the value of $2A - 3B - 2\sqrt{6}C$.

- (a) 25 (b) 7
 (c) 137 (d) 79

SSC CGL-(Tier-I) 16/08/2021 (Shift II)

Ans. (b) : From question,

$$\frac{(16\sqrt{2}x^3 + 81\sqrt{3}y^3)}{(2\sqrt{2}x + 3\sqrt{3}y)} = Ax^2 + By^2 + Cxy$$

From formula : $(a^3 + b^3) = (a + b)(a^2 + b^2 - ab)$
 $\frac{(2\sqrt{2}x + 3\sqrt{3}y)(8x^2 + 27y^2 - 6\sqrt{6}xy)}{(2\sqrt{2}x + 3\sqrt{3}y)} = Ax^2 + By^2 + Cxy$
 $\Rightarrow 8x^2 + 27y^2 - 6\sqrt{6}xy = Ax^2 + By^2 + Cxy$
 On comparing both sides,
 A = 8
 B = 27
 C = $-6\sqrt{6}$

Then,

$$\begin{aligned} 2A - 3B - 2\sqrt{6}C &= 2 \times 8 - 3 \times 27 - 2\sqrt{6} \times (-6\sqrt{6}) \\ &= 16 - 81 + 12 \times 6 \\ &= 16 - 81 + 72 \\ &= 88 - 81 \\ &= 7 \end{aligned}$$

$$\therefore \boxed{2A - 3B - 2\sqrt{6}C = 7}$$

67. If $4x^4 - 37x^2 + 9 = 0$, $x > \sqrt{\frac{3}{2}}$, then what is the value of $8x^3 - \frac{27}{x^3}$?

- (a) 35 (b) 215
(c) -215 (d) -35

SSC CGL-(Tier-I) 16/08/2021 (Shift II)

Ans. (b) : $4x^4 - 37x^2 + 9 = 0$

Let us consider $x^2 = a$

then, $4a^2 - 37a + 9 = 0$

$4a^2 - 36a - a + 9 = 0$

$4a(a-9) - 1(a-9) = 0$

$4a(a-9) - 1(a-9) = 0$

$a = 1/4$ or $a = 9$

On putting the value of $a = x^2$

$x^2 = 1/4$ not acceptable as $x > \sqrt{\frac{3}{2}}$

$x^2 = 9$ then $x = 3$ or $x = -3$

$$\left(x = -3 \text{ not valid as } x > \sqrt{\frac{3}{2}} \right)$$

Then put $x = 3$,

$$8x^3 - \frac{27}{x^3} = ?$$

$$? = 8 \times (3)^3 - \frac{27}{(3)^3}$$

$$? = 8 \times 27 - \frac{27}{27}$$

$$? = 216 - 1, \boxed{? = 215}$$

68. If $x + y + z = 1$, $xy + yz + zx = xyz = -4$, then what is the value of $(x^3 + y^3 + z^3)$?

- (a) 8 (b) -8
(c) 1 (d) -1

SSC CGL-(Tier-I) 18/08/2021 (Shift II)

Ans. (c) : Given,

$x + y + z = 1$, $xy + yz + zx = xyz = -4$

$(x^3 + y^3 + z^3) = ?$

$$\boxed{(x+y+z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)}$$

$$(1)^2 = x^2 + y^2 + z^2 + 2 \times (-4)$$

$$x^2 + y^2 + z^2 = 9$$

$$\therefore [x^3 + y^3 + z^3 - 3xyz = (x + y + z)[x^2 + y^2 + z^2 - xy - yz - zx]]$$

$$x^3 + y^3 + z^3 - 3 \times -4 = 1[9 - (-4)]$$

$$x^3 + y^3 + z^3 = 9 + 4 - 12$$

$$x^3 + y^3 + z^3 = 1$$

69. If $x + y = 2$ and $\frac{1}{x} + \frac{1}{y} = \frac{18}{5}$, then the value of $(x^3 + y^3)$ is :

- (a) $4\frac{2}{3}$ (b) $4\frac{3}{5}$
(c) $3\frac{1}{5}$ (d) $3\frac{1}{3}$

SSC CGL-(Tier-I) 16/08/2021 (Shift III)

Ans. (a) : Given:- $x + y = 2$ -----(i),

$$\frac{1}{x} + \frac{1}{y} = \frac{18}{5}$$

$$\frac{x+y}{xy} = \frac{18}{5}$$

On putting the value of $x + y$ from equation (i)

$$\frac{2}{xy} = \frac{18}{5}$$

$$xy = \frac{2 \times 5}{18} = \frac{5}{9} \text{ ----- (ii)}$$

On cubing of equation (i),

$$x^3 + y^3 + 3xy(x + y) = (2)^3$$

On putting the value of $xy = \frac{5}{9}$,

$$x^3 + y^3 + 3 \times \frac{5}{9} (2) = 8$$

$$x^3 + y^3 + \frac{10}{3} = 8$$

$$x^3 + y^3 = 8 - \frac{10}{3} = \frac{24 - 10}{3} = \frac{14}{3} = 4\frac{2}{3}$$

$$x^3 + y^3 = 4\frac{2}{3}$$

70. If $x - \frac{1}{x} = \sqrt{77}$, then one of the values of $x^3 + \frac{1}{x^3}$ is :

- (a) $80\sqrt{77}$ (b) 702
(c) $77\sqrt{77}$ (d) $3\sqrt{77}$

SSC CGL-(Tier-I) 18/08/2021 (Shift III)

Ans. (b) : Given :- $x - \frac{1}{x} = \sqrt{77}$

From formula $(a - b)^2 = (a + b)^2 - 4ab$

$$\left(\sqrt{77}\right)^2 = \left(x + \frac{1}{x}\right)^2 - 4$$

$$\left(x + \frac{1}{x}\right)^2 = 77 + 4 = 81$$

$$x + \frac{1}{x} = 9 \text{ -----(ii)}$$

On cubing both side of equation (ii),

$$\left(x + \frac{1}{x}\right)^3 = (9)^3$$

$$x^3 + \frac{1}{x^3} + 3 \times x \times \frac{1}{x} \left(x + \frac{1}{x} \right) = 729$$

$$x^3 + \frac{1}{x^3} + 3 \times (9) = 729$$

$$x^3 + \frac{1}{x^3} = 729 - 27 = 702$$

$$x^3 + \frac{1}{x^3} = 702$$

71. If $x + y + z = 3$, $xy + yz + zx = -12$ and $xyz = -16$, then the value of $\sqrt{x^3 + y^3 + z^3 + 13}$ is :
 (a) 9 (b) 8 (c) 10 (d) 11

SSC CGL-(Tier-I) 20/08/2021 (Shift III)

Ans. (c) : From formula :-

$$a^3 + b^3 + c^3 - 3abc = (a+b+c) [(a+b+c)^2 - 3(ab+bc+ca)]$$

$$x^3 + y^3 + z^3 - 3xyz = (x+y+z) [(x+y+z)^2 - 3(xy+yz+zx)]$$

As per question

$$x^3 + y^3 + z^3 - 3 \times (-16) = 3 [(3)^2 - 3(-12)]$$

$$x^3 + y^3 + z^3 + 48 = 3(9 + 36)$$

$$x^3 + y^3 + z^3 + 48 = 135$$

$$x^3 + y^3 + z^3 + 13 = 135 - 35 = 100$$

On taking square root both sides,

$$\sqrt{x^3 + y^3 + z^3 + 13} = \sqrt{100}$$

$$\sqrt{x^3 + y^3 + z^3 + 13} = 10$$

72. If $x^8 - 433x^4 + 16 = 0$, $x > 0$, then what is the value of $\left(x + \frac{2}{x}\right)$?

- (a) 7 (b) 4 (c) 5 (d) 9

SSC CGL-(Tier-I) 17/08/2021 (Shift II)

Ans. (c) : Given equation- $x^8 - 433x^4 + 16 = 0$

$$x^4 + \frac{16}{x^4} = 433$$

On adding +8 at both side

$$x^4 + \frac{16}{x^4} + 8 = 441$$

$$\left(x^2 + \frac{4}{x^2}\right)^2 = 21^2$$

$$x^2 + \frac{4}{x^2} = 21$$

$$x^2 + \frac{4}{x^2} + 4 = 21 + 4$$

$$\left(x + \frac{2}{x}\right)^2 = 25$$

$$x + \frac{2}{x} = 5$$

{ $\therefore x > 0$ }

73. If $(x+y)^3 + 27(x-y)^3 = (Ax-2y)(Bx^2+Cxy+13y^2)$, then the value of $A-B-C$ is :

- (a) 27 (b) 13
(c) 15 (d) 20

SSC CGL-(Tier-I) 17/08/2021 (Shift II)

Ans. (b) : Given that,

$$(x+y)^3 + 27(x-y)^3 = (Ax-2y)(Bx^2+Cxy+13y^2)$$

$$\text{LHS} = (x+y)^3 + (3x-3y)^3$$

$$(x+y+3x-3y)[(x+y)^2 + (3x-3y)^2 - (x+y)(3x-3y)]$$

$$\{ \therefore a^3 + b^3 = (a+b)(a^2 + b^2 - ab) \}$$

$$= (4x-2y)[10x^2 + 10y^2 - 16xy - 3(x^2 - y^2)]$$

$$= (4x-2y)[10x^2 + 10y^2 - 16xy - 3x^2 + 3y^2]$$

$$= (4x-2y)[7x^2 + 13y^2 - 16xy]$$

After comparing LHS with RHS,

$$(4x-2y)[7x^2 + 13y^2 - 16xy] = (Ax-2y)(Bx^2 + Cxy + 13y^2)$$

$$A = 4, B = 7 \text{ \& } C = -16$$

$$\therefore (A - B - C) = (4 - 7 + 16) = 13$$

74. If $\left(2x - \frac{3}{x}\right) = 2$, then what is the value of

$$\left(16x^4 + \frac{81}{x^4}\right) ?$$

- (a) 328 (b) 180
(c) 184 (d) 220

SSC CGL (Tier-I) 16/08/2021 (Shift I)

Ans. (c) : $\left(2x - \frac{3}{x}\right) = 2$

On squaring both sides,

$$4x^2 + \frac{9}{x^2} = 4 + 12$$

$$4x^2 + \frac{9}{x^2} = 16$$

Again on squaring both sides,

$$16x^4 + \frac{81}{x^4} = 256 - 72$$

$$16x^4 + \frac{81}{x^4} = 184$$

75. $x + y + z = 2$ and $xy + yz + zx = -11$, then the value of $x^3 + y^3 + z^3 - 3xyz$ is:

- (a) 78 (b) 71
(c) 74 (d) 69

SSC CGL (Tier-I) 16/08/2021 (Shift I)

Ans. (c) : $x + y + z = 2$

$$x^2 + y^2 + z^2 + 2(xy + yz + zx) = 4$$

$$x^2 + y^2 + z^2 = 4 + 22 = 26$$

$$\therefore x^3 + y^3 + z^3 - 3xyz = (x+y+z)[x^2 + y^2 + z^2 - (xy + yz + zx)]$$

$$= 2 \times [26 + 11]$$

$$= 2 \times 37 = 74$$

76. If $x + \frac{1}{x} = 4$, then the value of $x^5 + \frac{1}{x^5}$ is :

- (a) 776 (b) 684
(c) 724 (d) 736

SSC CGL-(Tier-I) 13/08/2021 (Shift I)

Ans. (c) : Given,

$$x + \frac{1}{x} = 4 \Rightarrow x^2 + \frac{1}{x^2} = 14, x^3 + \frac{1}{x^3} = 52$$

$$x^5 + \frac{1}{x^5} = \left(x^2 + \frac{1}{x^2}\right) \left(x^3 + \frac{1}{x^3}\right) - \left(x + \frac{1}{x}\right)$$

$$= 14 \times 52 - 4$$

$$= 728 - 4 = 724$$

77. If $x + y = 4$ and $\frac{1}{x} + \frac{1}{y} = \frac{16}{15}$, then what is the value of $(x^3 + y^3)$?

- (a) 18 (b) 16
(c) 19 (d) 21

SSC CGL-(Tier-I) 13/08/2021 (Shift I)

Ans. (c) :

Ans. (c) :

$$\frac{1}{x} + \frac{1}{y} = \frac{16}{15}$$

$$\frac{x+y}{xy} = \frac{16}{15}$$

$$\frac{4}{xy} = \frac{16}{15} \quad \dots(\text{Given } x + y = 4)$$

$$xy = \frac{15}{4}$$

$$\therefore x^3 + y^3 = (x+y)^3 - 3xy(x+y)$$

$$= 4^3 - 3 \times \frac{15}{4} \times 4$$

$$= 64 - 45 = 19$$

78. If $x + y + z = 3$, $x^2 + y^2 + z^2 = 45$ and $x^3 + y^3 + z^3 = 69$, then what is the value of xyz ?

- (a) -40 (b) 40
(c) -30 (d) 30

SSC CHSL 19/04/2021 (Shift-I)

Ans. (a) : Given,

$$x + y + z = 3, x^2 + y^2 + z^2 = 45, x^3 + y^3 + z^3 = 69, xyz = ?$$

$$(x+y+z)^2 = x^2 + y^2 + z^2 + 2(xy+yz+zx)$$

$$9 = 45 + 2(xy+yz+zx)$$

$$2(xy+yz+zx) = -36$$

$$xy+yz+zx = -18$$

$$x^3 + y^3 + z^3 - 3xyz = (x+y+z)(x^2 + y^2 + z^2 - xy - yz - zx)$$

$$69 - 3xyz = 3 \times [45 - (-18)]$$

$$-3xyz = 63 \times 3 - 69$$

$$xyz = \frac{189 - 69}{-3} = -40$$

79. If $x + \frac{1}{x} = \sqrt{7}$, then what is the value of $(x^2 + 1)$

$$\div \left[x^4 + \left(\frac{1}{x^2} \right) \right] ?$$

- (a) $2\sqrt{7}$ (b) $3\sqrt{7}$
(c) $\frac{1}{2}$ (d) $\frac{1}{4}$

SSC CHSL 19/04/2021 (Shift-I)

Ans. (d) : Given,

$$x + \frac{1}{x} = \sqrt{7}$$

$$x^3 + \frac{1}{x^3} = (\sqrt{7})^3 - 3\sqrt{7} = 4\sqrt{7}$$

$$\frac{x^2 + 1}{x^4 + \frac{1}{x^2}} = ?$$

On dividing by x ,

$$? = \frac{x + \frac{1}{x}}{x^3 + \frac{1}{x^3}}$$

$$? = \frac{\sqrt{7}}{4\sqrt{7}} = \frac{1}{4}$$

80. If $x^6 - 6\sqrt{6}y^6 = (x^2 + Ay^2)(x^4 + Bx^2y^2 + Cy^4)$, then what will be the value of $(A^2 - B^2 + C^2)$?

- (a) 27 (b) 42
(c) 36 (d) 18

SSC CHSL 10/08/2021 (Shift-I)

Ans. (c) : $x^6 - 6\sqrt{6}y^6 = (x^2 + Ay^2)(x^4 + Bx^2y^2 + Cy^4)$

$$(x^2)^3 - (\sqrt{6}y^2)^3 = (x^2 + Ay^2)(x^4 + Bx^2y^2 + Cy^4)$$

$$(x^2 - \sqrt{6}y^2) [x^4 + \sqrt{6}x^2y^2 + 6y^4]$$

$$= (x^2 + Ay^2)(x^4 + Bx^2y^2 + Cy^4)$$

On comparing both sides,

$$\therefore A = -\sqrt{6}, \quad B = \sqrt{6}, \quad C = 6$$

$$\therefore A^2 - B^2 + C^2 = (-\sqrt{6})^2 - (\sqrt{6})^2 + (6)^2$$

$$= 6 - 6 + 36 = 36$$

81. If $x + \frac{1}{15x} = 3$, then the value of $9x^3 + \frac{1}{375x^3}$

will be:

- (a) 237.6 (b) 376.2
(c) 273.6 (d) 367.2

SSC CHSL 10/08/2021 (Shift-I)

Ans. (a) : $x + \frac{1}{15x} = 3$

On multiplying by 3 in both side,

$$3x + \frac{1}{5x} = 9$$

Taking cube on both sides,

$$27x^3 + \frac{1}{125x^3} + 3 \times 3x \times \frac{1}{5x} \left(3x + \frac{1}{5x} \right) = 729$$

$$27x^3 + \frac{1}{125x^3} + \frac{9}{5} \times 9 = 729$$

On multiplying by $\frac{1}{3}$ of both side

$$9x^3 + \frac{1}{375x^3} + 5.4 = 243$$

$$9x^3 + \frac{1}{375x^3} = 237.6$$

82. If $x - y = 4$ and $xy = 3$, then what is the value of $x^3 - y^3$?

- (a) 88 (b) 28
(c) 100 (d) 64

SSC CHSL 06/08/2021 (Shift-I)

Ans. (c) : $x - y = 4$, $xy = 3$
 $\therefore x^3 - y^3 = (x-y)(x^2 + y^2 + xy)$ $[(x^2 + y^2 = (x-y)^2 + 2xy)]$
 $= 4 \times [(x-y)^2 + 2xy + xy]$
 $= 4 \times [4^2 + 3 \times 3]$
 $= 4 \times [16 + 9]$
 $= 4 \times 25 = 100$

83. If $x^2 - 6\sqrt{3}x + 1 = 0$, then the value of $x^3 + \frac{1}{x^3}$ will be:

- (a) $666\sqrt{3}$ (b) $630\sqrt{3}$
(c) $234\sqrt{3}$ (d) $216\sqrt{3}$

SSC CHSL 12/04/2021 (Shift-I)

Ans : (b) $x^2 - 6\sqrt{3}x + 1 = 0$ ----- [Given]

$$x - 6\sqrt{3} + \frac{1}{x} = 0$$

$$x + \frac{1}{x} = 6\sqrt{3}$$

On cubing both sides,

$$x^3 + \frac{1}{x^3} + 3\left(x + \frac{1}{x}\right) = 6 \times 6 \times 6 \times 3\sqrt{3}$$

$$x^3 + \frac{1}{x^3} = 648\sqrt{3} - 3 \times 6\sqrt{3}$$

$$= 630\sqrt{3}$$

84. If $a + b = p$, $ab = q$, then $(a^4 + b^4)$ is equal to

- (a) $p^4 - 2p^2q^2 + q^2$ (b) $p^4 - 4p^2q^2 + 2q^2$
(c) $p^4 - 4p^2q + q^2$ (d) $p^4 - 4p^2q + 2q^2$

SSC CHSL 04/08/2021 (Shift-I)

Ans. (d) : Given—
 $a + b = p$, $ab = q$ $[a^4 + b^4 = ?]$
 $(a + b) = p$ (On squaring both sides)
 $a^2 + b^2 + 2ab = p^2$
 $a^2 + b^2 = p^2 - 2q$ (On squaring both sides)
 $a^4 + b^4 + 2q^2 = p^4 + 4q^2 - 2p^2 \times 2q$
 $a^4 + b^4 = p^4 - 4p^2q + 2q^2$

85. If $x^4 + y^4 + x^2y^2 = 117$ and $x^2 + y^2 - xy = 3(4 + \sqrt{3})$, then the value of $(x^2 + y^2)$ will be:

- (a) $6\sqrt{3}$ (b) 12
(c) 9 (d) $13\sqrt{3}$

SSC CHSL 12/04/2021 (Shift-I)

Ans : (b) $x^4 + y^4 + x^2y^2 = 117$, $x^2 + y^2 - xy = 3(4 + \sqrt{3})$
 $x^4 + y^4 + x^2y^2 = (x^2 + y^2 + xy)(x^2 + y^2 - xy)$

$$117 = (x^2 + y^2 + xy) [3(4 + \sqrt{3})]$$

$$x^2 + y^2 + xy = \frac{117}{3(4 + \sqrt{3})} \times \frac{(4 - \sqrt{3})}{(4 - \sqrt{3})}$$

$$= \frac{39(4 - \sqrt{3})}{13} = 3(4 - \sqrt{3})$$

$$x^2 + y^2 + xy = 3(4 - \sqrt{3}) \quad \dots(i)$$

$$x^2 + y^2 - xy = 3(4 + \sqrt{3}) \quad \dots(ii)$$

From eqⁿ (i) and eqⁿ (ii),

$$2(x^2 + y^2) = 24$$

$$x^2 + y^2 = 12$$

86. If $\left(x + \frac{1}{x}\right)^2 = 27$, then what is the value of

$\left(x^2 + \frac{1}{x^2}\right)$? Given that x is real.

- (a) 11 (b) 25
(c) 7 (d) 9

SSC CHSL 04/08/2021 (Shift-I)

Ans. (b) $\left(x + \frac{1}{x}\right)^2 = 27$

$$x^2 + \frac{1}{x^2} + 2 \times x \times \frac{1}{x} = 27$$

$$x^2 + \frac{1}{x^2} = 27 - 2$$

$$x^2 + \frac{1}{x^2} = 25$$

87. If $\left(x + \frac{2}{x}\right) = 7$, then what is the value of

$\left(2x^2 + \frac{8}{x^2}\right)$?

- (a) 90 (b) 44
(c) 50 (d) 94

SSC CHSL 16/04/2021 (Shift-I)

Ans. (a) : Given, $x + \frac{2}{x} = 7$

On squaring both sides,

$$x^2 + \frac{4}{x^2} + 4 = 49$$

$$x^2 + \frac{4}{x^2} = 45$$

On multiplying by 2 on both sides,

$$2x^2 + \frac{8}{x^2} = 90$$

88. If $x - 3 = \frac{1}{2x}$, then what is the value of

$\left(x^4 + \frac{1}{16x^4}\right)$?

- (a) 11 (b) $99\frac{1}{2}$
 (c) 98 (d) 10

SSC CHSL 16/04/2021 (Shift-I)

Ans. (b) : $x - 3 = \frac{1}{2x}$

$$x - \frac{1}{2x} = 3$$

On squaring both sides,

$$x^2 + \frac{1}{4x^2} - 2 \times x \times \frac{1}{2x} = 9$$

$$x^2 + \frac{1}{4x^2} = 10$$

Again on squaring both sides,

$$x^4 + \frac{1}{16x^4} + 2 \times x^2 \times \frac{1}{4x^2} = 100$$

$$x^4 + \frac{1}{16x^4} = 100 - \frac{1}{2} = \frac{199}{2} = 99\frac{1}{2}$$

89. Given that $3\sqrt{3}x^3 - 8y^3 = (\sqrt{3}x + Ay)(3x^2 + By^2 + Cxy)$, the value of $(A^2 + B^2 - C^2)$ is:
 (a) 0 (b) 12
 (c) 8 (d) 4

SSC CHSL 12/08/2021 (Shift-I)

Ans. (c) : $3\sqrt{3}x^3 - 8y^3 = (\sqrt{3}x + Ay)(3x^2 + By^2 + Cxy)$
 $(\sqrt{3}x - 2y)(3x^2 + 2\sqrt{3}xy + 4y^2)$
 $= (\sqrt{3}x + Ay)(3x^2 + By^2 + Cxy)$

$$\therefore a^3 - b^3 = (a-b)(a^2 + ab + b^2)$$

\therefore On comparing both sides,

$$A = -2, B = 4, C = 2\sqrt{3}$$

So, $A^2 + B^2 - C^2 = (-2)^2 + (4)^2 - (2\sqrt{3})^2$
 $= 4 + 16 - 12 = 8$

90. If $3x - 2y + 3 = 0$, then what will be the value of $27x^3 + 54xy + 30 - 8y^3$?
 (a) 3 (b) -27
 (c) -57 (d) 57

SSC CHSL 12/08/2021 (Shift-I)

Ans. (a) : $3x - 2y + 3 = 0$

Let, $x = 1, y = 3$

Then, $3x - 2y + 3 = 3 \times 1 - 2 \times 3 + 3 = 0$ L.H.S = R.H.S.

On putting $x = 1$ and $y = 3$,

$$27x^3 + 54xy + 30 - 8y^3 = 27 + 54 \times 1 \times 3 + 30 - 8 \times 27$$

$$= 27 + 162 + 30 - 216$$

$$= 219 - 216 = 3$$

91. If $\sqrt{x} + \frac{1}{\sqrt{x}} = 2\sqrt{3}$, then what will be the value of $x^4 + \frac{1}{x^4}$?

- (a) 10406 (b) 9602
 (c) 9606 (d) 10402

SSC CHSL 12/08/2021 (Shift-I)

Ans. (b) : $\sqrt{x} + \frac{1}{\sqrt{x}} = 2\sqrt{3}$

On squaring both sides,

$$x + \frac{1}{x} + 2 = 12$$

$$x + \frac{1}{x} = 10$$

Again, on squaring both sides,

$$x^2 + \frac{1}{x^2} = 98$$

Again on squaring both sides,

$$x^4 + \frac{1}{x^4} = (98)^2 - 2 = 9604 - 2 = 9602$$

92. If $3u + 2v = 7$ and $uv = 2$, then the value of $(3u - 2v)$ is:

- (a) 2 (b) 0
 (c) 1 (d) 5

SSC CHSL 13/04/2021 (Shift-I)

Ans. (c) : $\therefore (a-b)^2 = (a+b)^2 - 4ab$
 $(3u - 2v)^2 = (3u + 2v)^2 - 4 \times 3u \times 2v$
 $= 49 - 24 \times 2 = 1$
 $3u - 2v = 1$

93. If $2x^2 - 6x = 1$, then $x^2 + \frac{1}{4x^2} = ?$
 (a) 8 (b) 12
 (c) 9 (d) 10

SSC CHSL 13/04/2021 (Shift-I)

Ans. (d) : $2x^2 - 6x = 1$

On dividing by $2x$,

$$x - 3 = \frac{1}{2x}$$

$$x - \frac{1}{2x} = 3$$

On squaring both sides,

$$\left(x - \frac{1}{2x}\right)^2 = 9$$

$$x^2 + \frac{1}{4x^2} - 2 \times x \times \frac{1}{2x} = 9$$

$$x^2 + \frac{1}{4x^2} = 10$$

94. If $a + b + c = 5$ and $a^3 + b^3 + c^3 - 3abc = 185$, then the value of $ab + bc + ca$ lies between:

- (a) -7 and -3 (b) 1 and 5
 (c) -3 and 1 (d) 5 and 9

SSC CHSL 13/04/2021 (Shift-I)

Ans. (a) :
 $a^3 + b^3 + c^3 - 3abc = (a+b+c) [(a+b+c)^2 - 3(ab+bc+ca)]$
 $185 = 5[25 - 3(ab+bc+ca)]$
 $37 - 25 = -3(ab+bc+ca)$
 $ab + bc + ca = -4$

$\therefore -4$ lies between -7 and -3 .

Hence option (a) will be right.

95. If $3x + 5y = 14$ and $xy = 6$, then what is the value of $9x^2 + 25y^2$?

- (a) 16 (b) 14
(c) 20 (d) 182

SSC CHSL 11/08/2021 (Shift-I)

Ans. (a) : $3x + 5y = 14$, $xy = 6$ (Given)

On squaring both sides,

$$(3x + 5y)^2 = 14^2$$

$$9x^2 + 25y^2 + 30xy = 196 \quad (\because xy = 6)$$

So, $9x^2 + 25y^2 = 196 - 30 \times 6 = 196 - 180 = 16$

96. If $a^2 + b^2 + c^2 + 48 = 8(a + b + c)$, then what is the value of $\sqrt[3]{a^3 - b^3 + c^3}$?

- (a) 6 (b) 4
(c) 3 (d) 2

SSC CHSL 11/08/2021 (Shift-I)

Ans. (b) : $a^2 + b^2 + c^2 + 48 = 8(a + b + c)$

$$a^2 - 8a + 16 + b^2 - 8b + 16 + c^2 - 8c + 16 = 0$$

$$(a-4)^2 + (b-4)^2 + (c-4)^2 = 0$$

$$\therefore (a-4)^2 = 0, \quad a = 4$$

$$(b-4)^2 = 0, \quad b = 4$$

$$(c-4)^2 = 0, \quad c = 4$$

So, $\sqrt[3]{a^3 - b^3 + c^3} = \sqrt[3]{4^3 - 4^3 + 4^3} = 4$

97. If $x^4 + x^{-4} = 47$, $x > 0$, then the value of $(2x - 3)^2$ is:

- (a) 9 (b) 3
(c) 5 (d) 7

SSC CHSL 11/08/2021 (Shift-I)

Ans. (c) : $x^4 + \frac{1}{x^4} = 47$

$$x^2 + \frac{1}{x^2} = 7$$

$$\left(x + \frac{1}{x}\right)^2 = 7 + 2 = 9$$

$$x + \frac{1}{x} = 3$$

$$x^2 + 1 - 3x = 0$$

$$x^2 - 3x + 1 = 0$$

Multiply by 4 on both sides,

$$4x^2 - 12x + 4 = 0$$

After adding 5 on both sides,

$$4x^2 - 12x + 4 + 5 = 5$$

$$4x^2 - 12x + 9 = 5$$

$$(2x - 3)^2 = 5$$

98. If $x - \frac{2}{x} = 4$, then what will be the value of

$$x^2 + \frac{4}{x^2}?$$

- (a) 8 (b) 20
(c) 18 (d) 12

SSC CHSL 04/08/2021 (Shift-II)

Ans. (b) : $x - \frac{2}{x} = 4$

$$x^2 + \frac{4}{x^2} - 2 \times x \times \frac{2}{x} = 16 \text{ (on squaring both sides)}$$

$$x^2 + \frac{4}{x^2} = 16 + 4 = 20$$

99. If $\sqrt{x} + \frac{1}{\sqrt{x}} = \sqrt{6}$, then the value of $x^6 + \frac{1}{x^6}$ will

be:

- (a) 2712 (b) 2270
(c) 2502 (d) 2702

SSC CHSL 04/08/2021 (Shift-II)

Ans. (d) : $\sqrt{x} + \frac{1}{\sqrt{x}} = \sqrt{6}$

On squaring both sides,

$$x + \frac{1}{x} + 2 = 6$$

$$x + \frac{1}{x} = 4$$

$$x^2 + \frac{1}{x^2} = 14$$

Again on cubing both sides,

$$x^6 + \frac{1}{x^6} = (14)^3 - 3 \times 14$$

$$= 2744 - 42 = 2702$$

100. If $x^4 + \frac{1}{x^4} = 3842$, then the positive value of

$x + \frac{1}{x}$ will be:

- (a) 10 (b) 8
(c) 12 (d) 6

SSC CHSL 04/08/2021 (Shift-II)

Ans. (b) : $x^4 + \frac{1}{x^4} = 3842$

$$x^2 + \frac{1}{x^2} = \sqrt{3842 + 2} = \sqrt{3844} = 62$$

$$x + \frac{1}{x} = \sqrt{62 + 2} = \sqrt{64} = 8$$

Hence, $x + \frac{1}{x} = 8$

101. If $x + \frac{1}{3x} = 5$, then the value of $27x^3 + \frac{1}{x^3}$ will

be:

- (a) 3042 (b) 3024
(c) 3420 (d) 3240

SSC CHSL 10/08/2021 (Shift-II)

Ans. (d) : Given

$$x + \frac{1}{3x} = 5$$

On multiplying by 3 of both side

$$3x + \frac{1}{x} = 5 \times 3 = 15$$

On cubing both sides,

$$\left(3x + \frac{1}{x}\right)^3 = (15)^3$$

$$27x^3 + \frac{1}{x^3} + 3 \times 3x \times \frac{1}{x} \left(3x + \frac{1}{x}\right) = 3375$$

$$27x^3 + \frac{1}{x^3} + 9(15) = 3375$$

$$27x^3 + \frac{1}{x^3} = 3375 - 135$$

$$27x^3 + \frac{1}{x^3} = 3240$$

102. If $1 + 4x^2 + 16x^4 = 512$, and $1 - 2x + 4x^2 = 64$, then the value of $1 - 2x + 4x^2$ is:

- (a) 6 (b) 8
(c) 10 (d) 12

SSC CHSL 10/082021 (Shift-II)

Ans. (b) : Given :-

$$1 + 4x^2 + 16x^4 = 512 \text{ -----(i)}$$

$$\text{And } 1 - 2x + 4x^2 = 64 \text{ ----- (ii)}$$

On dividing equation (i) by equation (ii)

$$\frac{1 + 4x^2 + 16x^4}{1 - 2x + 4x^2} = \frac{512}{64}$$

$$\frac{(1 + 2x + 4x^2)(1 - 2x + 4x^2)}{1 - 2x + 4x^2} = 8$$

$$1 + 2x + 4x^2 = 8$$

103. If $x^4 - 12x^2 + 1 = 0$, then what will be the value of $x^4 + \frac{1}{x^4}$?

- (a) 142 (b) 146
(c) 10 (d) 144

SSC CHSL 06/08/2021 (Shift-III)

Ans. (a) : Given :- $x^4 - 12x^2 + 1 = 0$

$$x^4 + 1 = 12x^2$$

On dividing by x^2 of both side

$$x^2 + \frac{1}{x^2} = 12$$

On squaring both side

$$\left(x^2 + \frac{1}{x^2}\right)^2 = (12)^2$$

$$x^4 + \frac{1}{x^4} + 2 = 144$$

$$x^4 + \frac{1}{x^4} = 144 - 2$$

$$x^4 + \frac{1}{x^4} = 142$$

104. If $x + \frac{81}{x} = 18$ where $x > 0$, then the value of

$x^2 + \frac{162}{x^2}$ is:

- (a) 78 (b) 83
(c) 85 (d) 81

SSC CHSL 06/08/2021 (Shift-III)

Ans. (b) : From question,

$$x + \frac{81}{x} = 18$$

$$\Rightarrow x^2 + 81 = 81x$$

$$\Rightarrow x^2 - 18x + 81 = 0$$

$$\Rightarrow x^2 - (9+9)x + 81 = 0$$

$$\Rightarrow x^2 - 9x - 9x + 81 = 0$$

$$\Rightarrow x(x-9) - 9(x-9) = 0$$

$$\Rightarrow (x-9)(x-9) = 0$$

$$\therefore x = 9$$

$$\text{So, } x^2 + \frac{162}{x^2} = 9^2 + \frac{162}{9^2}$$

$$= 81 + \frac{162}{81}$$

$$= 81 + 2$$

$$= 83$$

105. If $x^3 + y^3 = 468$ and $x + y = 12$, then the value of $x^4 + y^4$ will be:

- (a) 3026 (b) 2036
(c) 3620 (d) 3025

SSC CHSL 13/04/2021 (Shift-III)

Ans.(a) : Given :- $x^3 + y^3 = 468$

$$(x + y)(x^2 + y^2 - xy) = 468$$

$$x^2 + y^2 - xy = \frac{468}{12} \text{ ----- } [\because \text{on putting the value of } (x+y) = 12]$$

$$x^2 + y^2 - xy = 39 \text{ -----(i)}$$

$$\text{And } x + y = 12$$

On squaring both sides

$$x^2 + y^2 + 2xy = 144 \text{ -----(ii)}$$

On subtracting equation (i) from equation (ii)

$$xy = 144 - 39 = \frac{105}{3} = 35 \text{ -----(iii)}$$

On putting the value of xy in equation (ii)

$$x^2 + y^2 + 2 \times 35 = 144$$

$$x^2 + y^2 = 144 - 70 = 74$$

Again, on squaring both sides,

$$x^4 + y^4 + 2x^2y^2 = (74)^2$$

On putting the value of $xy = 35$,

$$x^4 + y^4 + 2 \times (35)^2 = 5476$$

$$x^4 + y^4 = 5476 - 2450 = 3026$$

Hence, $x^4 + y^4 = 3026$

106. If $x^2 - 3\sqrt{2}x + 1 = 0$, then what is the value of $x^3 + \left(\frac{1}{x^2}\right)$?

- (a) $30\sqrt{6}$ (b) $45\sqrt{2}$
 (c) $15\sqrt{6}$ (d) $30\sqrt{2}$

SSC CHSL 04/08/2021 (Shift-III)

Ans. (b) : $x^2 - 3\sqrt{2}x + 1 = 0$

$$x^2 + 1 = 3\sqrt{2}x$$

On dividing by x both sides,

$$x + \frac{1}{x} = 3\sqrt{2}$$

On cubing both sides,

$$x^3 + \frac{1}{x^3} + 3\left(x + \frac{1}{x}\right) = 54\sqrt{2}$$

$$x^3 + \frac{1}{x^3} + 3 \times 3\sqrt{2} = 54\sqrt{2}$$

$$x^3 + \frac{1}{x^3} = 54\sqrt{2} - 9\sqrt{2}$$

$$x^3 + \frac{1}{x^3} = 45\sqrt{2}$$

107. If $x^2 + 1 - 2x = 0$, $x > 0$, then $x^2(x^2 - 2) = \dots\dots\dots$

- (a) 1 (b) $\sqrt{2}$
 (c) -1 (d) 0

SSC CHSL 04/08/2021 (Shift-III)

Ans. (c) : $x^2 + 1 - 2x = 0$ ($x > 0$)

$$(x-1)^2 = 0$$

$$x = 1$$

$$x^2(x^2 - 2) = 1(1 - 2) = -1$$

108. If $a + b = 24$ and $a^2 + b^2 = 306$, where $a > b$, then the value of $4a - 5b$ is:

- (a) 18 (b) 20
 (c) 12 (d) 15

SSC CHSL 05/08/2021 (Shift-III)

Ans. (d) : $(a + b) = 24$ -----(i) [Given]

On squaring both sides,

$$a^2 + b^2 + 2ab = 576$$

$$2ab = 576 - (a^2 + b^2)$$

$$2ab = 576 - 306 = 270$$

$$ab = \frac{270}{2} = 135$$

$$ab = 135 \dots\dots(ii)$$

From equations (i) and (ii)

$$a = 15, b = 9$$

Then, $4a - 5b = 4 \times 15 - 5 \times 9 = 15$, $4a - 5b = 15$

109. If $x - y = 4$ and $x^3 - y^3 = 316$, then the value of $x^4 + y^4$ is:

- (a) 2248 (b) 2482
 (c) 2428 (d) 2284

SSC CHSL 05/08/2021 (Shift-III)

Ans. (b) : Given :- $x - y = 4$ and $x^3 - y^3 = 316$

From, $x^3 - y^3 = 316$

$$(x - y)(x^2 + y^2 + xy) = 316$$

$$x^2 + y^2 + xy = \frac{316}{4} = 79 \dots\dots\dots(i)$$

From, $x - y = 4$

On squaring both sides,

$$x^2 + y^2 - 2xy = 16 \dots\dots\dots(ii)$$

From equation (i) and (ii),

$$xy = 21 = 7 \times 3$$

On taking $x = 7$ and $y = 3$,

$$\text{Now, } x^4 + y^4 = (7)^4 + (3)^4 = 2401 + 81$$

$$x^4 + y^4 = 2482$$

109. If $a + b + c = 5$, $a^2 + b^2 + c^2 = 27$ and $a^3 + b^3 + c^3 = 125$ then the value of $\frac{abc}{5}$ is:

125 then the value of $\frac{abc}{5}$ is:

- (a) -1 (b) -5
 (c) 1 (d) 5

SSC CHSL 10/08/2021 (Shift-III)

Ans. (a) : Given :-

$$a + b + c = 5, a^2 + b^2 + c^2 = 27 \text{ and } a^3 + b^3 + c^3 = 125$$

On squaring both sides of $a + b + c = 5$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = 25$$

$$2(ab + bc + ca) = 25 - 27 = -2$$

$$ab + bc + ca = \frac{-2}{2} = -1 \dots\dots(i)$$

Now from formula :

$$a^3 + b^3 + c^3 - 3abc = (a + b + c) [(a+b+c)^2 - 3(ab+bc+ca)]$$

As per question

$$a^3 + b^3 + c^3 - 3abc = 5 [(5)^2 - 3(-1)]$$

$$125 - 3abc = 5(25 + 3)$$

$$3abc = 125 - 140$$

$$abc = \frac{-15}{3} = -5$$

On dividing by 5 both sides,

$$\frac{abc}{5} = \frac{-5}{5} = -1$$

$$\frac{abc}{5} = -1$$

110. If $a + b + c = 11$ and $ab + bc + ca = 15$ then what is the value of $a^3 + b^3 + c^3 - 3abc$?

- (a) 368 (b) 386
 (c) 638 (d) 836

SSC CHSL 10/08/2021 (Shift-III)

Ans. (d) : From formula :-

$$a^3 + b^3 + c^3 - 3abc = (a+b+c) [(a+b+c)^2 - 3(ab+bc+ca)]$$

$$a^3 + b^3 + c^3 - 3abc = 11 [(11)^2 - 3(15)]$$

$$= 11(121 - 45)$$

$$a^3 + b^3 + c^3 - 3abc = 11 \times 76 = 836$$

Hence, $a^3 + b^3 + c^3 - 3abc = 836$

111. If $(x-1.5)^3 + (x-4)^3 + (x-3.5)^3 = 3(x-1.5)(x-4)(x-3.5)$, then what is the value of x?

- (a) 9 (b) 3
 (c) 6 (d) 1

SSC CHSL 10/08/2021 (Shift-III)

Ans. (b) :

$$(x-1.5)^3 + (x-4)^3 + (x-3.5)^3 = 3(x-1.5)(x-4)(x-3.5)$$

Now from the formula

$$\text{When, } a^3 + b^3 + c^3 = 3abc$$

$$\text{Then, } a + b + c = 0$$

$$\text{Hence } x-1.5 + x-4 + x-3.5 = 0$$

$$3x - 9 = 0$$

$$3x = 9$$

$$x = \frac{9}{3} = 3$$

112. If $a^3 + b^3 + c^3 - 3abc = 250$ and $a + b + c = 10$, then what will be the value of $\frac{1}{5}(ab + bc + ca)$?

(a) 10

(b) 25

(c) 15

(d) 5

SSC CHSL 16/04/2021 (Shift-III)

Ans.(d) : Given :- $a^3 + b^3 + c^3 - 3abc = 250$

$$\text{and } a+b+c = 10$$

From formula,

$$a^3 + b^3 + c^3 - 3abc = (a+b+c) [(a+b+c)^2 - 3(ab+bc+ca)]$$

$$250 = 10 [(10)^2 - 3(ab+bc+ca)]$$

$$100 - 3(ab+bc+ca) = \frac{250}{10} = 25$$

$$3(ab+bc+ca) = 100 - 25 = 75$$

$$ab+bc+ca = \frac{75}{3} = 25$$

Hence

$$\frac{1}{5}(ab+bc+ca) = \frac{1}{5} \times 25 = 5$$

113. If $x^2 + y^2 = 45$ and $x-y = 5$ then what is the value of $x^3 - y^3$?

(a) -25

(b) 250

(c) 275

(d) 150

SSC CHSL 16/04/2021 (Shift-III)

Ans.(c) : Given :-

$$x-y=5$$

On squaring both sides,

$$x^2 + y^2 - 2xy = 25$$

On putting the value of $x^2 + y^2 = 45$

$$2xy = 45 - 25 = 20$$

$$xy = \frac{20}{2} = 10$$

$$\text{Now } x^3 - y^3 = (x-y)(x^2 + y^2 + xy)$$

$$= 5(45+10)$$

$$= 5 \times 55 = 275$$

$$\therefore x^3 - y^3 = 275$$

114. If $a^2 + 49b^2 + c^2 + 18 = 2(28b - c - a)$ then the value of $(a + 7b - c)$ is:

(a) 4

(b) 2

(c) -1

(d) 6

SSC CHSL 19/04/2021 (Shift-III)

Ans. (a) : Given :-

$$a^2 + 49b^2 + c^2 + 18 = 2(28b - c - a)$$

$$a^2 + 2a + 49b^2 - 56b + c^2 + 2c + 18 = 0$$

$$a^2 + 2a + 1 + (7b)^2 - 56b + 16 + c^2 + 2c + 1 = 0$$

$$(a+1)^2 + (7b-4)^2 + (c+1)^2 = 0$$

Hence,

$$a = -1$$

$$b = \frac{4}{7},$$

and $c = -1$

On putting the value of a, b and c in $a + 7b - c$,

$$-1 + 7 \times \frac{4}{7} + 1$$

$$-1 + 4 + 1 = 4$$

Hence,

$$\boxed{a + 7b - c = 4}$$

115. If $x - y - z = 0$, then the value of $(x^2 + y^2 + z^2) \div (y^2 + xz)$ is:

(a) -1

(b) 2

(c) 1

(d) -2

SSC CHSL 12/04/2021 (Shift-III)

Ans : (b) Given :- $x - y - z = 0$

On taking, $x = 2, y = 1$ and $z = 1$

$$x - y - z = 2 - 1 - 1 = 0$$

Hence,

$$(x^2 + y^2 + z^2) \div (y^2 + xz)$$

$$[(2)^2 + (1)^2 + (1)^2] \div [(1)^2 + 2 \times 1]$$

$$6 \div 3 = 2$$

$$(x^2 + y^2 + z^2) \div (y^2 + xy) = 2$$

116. If $x^4 + \frac{1}{x^4} = 6887$, then the positive value of $x - \frac{1}{x}$ is ?

(a) 9

(b) 8

(c) 12

(d) 15

SSC CHSL 12/04/2021 (Shift-III)

Ans : (a) Given :-

$$x^4 + \frac{1}{x^4} = 6887$$

On adding 2 at both sides,

$$x^4 + \frac{1}{x^4} + 2 = 6887 + 2 = 6889$$

$$\left(x^2 + \frac{1}{x^2}\right)^2 = (83)^2$$

$$x^2 + \frac{1}{x^2} = 83$$

On subtracting 2 from both sides,

$$x^2 + \frac{1}{x^2} - 2 = 83 - 2 = 81$$

$$\left(x - \frac{1}{x}\right)^2 = (9)^2$$

$$x - \frac{1}{x} = 9$$

117. If $x^2 - 3x + 1 = 0$, then the value of

$$2\left(x^8 + \frac{1}{x^8}\right) - 5\left(x^2 + \frac{1}{x^2}\right) \text{ is:}$$

- (a) 4370 (b) 4279
(c) 4379 (d) 3479

SSC CHSL 12/04/2021 (Shift-III)

Ans : (c) Given :- $x^2 - 3x + 1 = 0$

$$x + \frac{1}{x} = 3$$

On squaring both sides,

$$x^2 + \frac{1}{x^2} = 9 - 2 = 7 \text{ -----(ii)}$$

Again, on squaring both sides,

$$x^4 + \frac{1}{x^4} = 49 - 2 = 47$$

Again, on squaring both sides,

$$x^8 + \frac{1}{x^8} = 2209 - 2 = 2207 \text{ -----(iii)}$$

From equation (ii) and (iii),

$$2\left(x^8 + \frac{1}{x^8}\right) - 5\left(x^2 + \frac{1}{x^2}\right) = 4414 - 35 = 4379$$

Hence, $2\left(x^8 + \frac{1}{x^8}\right) - 5\left(x^2 + \frac{1}{x^2}\right) = 4379$

118. If $(4x - 5)^3 + (x - 2)^3 + 27(2x - 5)^3 = 9(4x - 5)(x - 2)(2x - 5)$, then the value of $\left(x + \frac{3}{2}\right)$ will be:

- (a) $\frac{1}{2}$ (b) $\frac{3}{2}$
(c) $\frac{7}{2}$ (d) $\frac{5}{2}$

SSC CHSL 05/08/2021 (Shift-II)

Ans. (c) : Given :-

$$(4x-5)^3 + (x-2)^3 + 27(2x-5)^3 = 9(4x-5)(x-2)(2x-5)$$

Now, from formula

$$\text{When } a^3 + b^3 + c^3 = 3abc$$

$$\text{Then } a+b+c=0$$

$$\text{Hence, } 4x-5 + x-2 + 3(2x-5) = 0$$

$$4x-5 + x-2 + 6x-15 = 0$$

$$11x - 22 = 0$$

$$11x = 22$$

$$x = \frac{22}{11} = 2$$

$$\left(x + \frac{3}{2}\right) = 2 + \frac{3}{2} = \frac{7}{2}$$

119. If $x^2 - 5\sqrt{2}x - 1 = 0$, then what will be the value of $x^3 - \frac{1}{x^3}$?

- (a) $250\sqrt{2}$ (b) $485\sqrt{2}$
(c) $265\sqrt{2}$ (d) $255\sqrt{2}$

SSC CHSL 19/08/2021 (Shift-II)

Ans. (c) : Given :-

$$x^2 - 5\sqrt{2}x - 1 = 0$$

$$x^2 - 1 = 5\sqrt{2}x$$

$$x - \frac{1}{x} = 5\sqrt{2}$$

On cubing both sides,

$$x^3 - \frac{1}{x^3} - 3\left(x - \frac{1}{x}\right) = 250\sqrt{2}$$

$$x^3 - \frac{1}{x^3} - 3 \times 5\sqrt{2} = 250\sqrt{2}$$

$$x^3 - \frac{1}{x^3} - 15\sqrt{2} = 250\sqrt{2}$$

$$x^3 - \frac{1}{x^3} = 265\sqrt{2}$$

120. If $a^4 + b^4 + a^2b^2 = 133$ and $a^2 + b^2 - ab = 19$, then the value of ab will be:

- (a) -9 (b) 15
(c) -6 (d) 12

SSC CHSL 19/08/2021 (Shift-II)

Ans. (c) : From the formula,

$$a^4 + b^4 + a^2b^2 = (a^2 + b^2 - ab)(a^2 + b^2 + ab)$$

As per question:-

$$133 = 19(a^2 + b^2 + ab)$$

$$a^2 + b^2 + ab = \frac{133}{19} = 7 \text{ -----(i)}$$

and $a^2 + b^2 - ab = 19$ ----- (ii) -----(Given)

On solving equation (i) and (ii),

$$2ab = -12$$

$$ab = \frac{-12}{2} = -6$$

121. If $x - y = \frac{7}{4}$ and $\frac{1}{x} - \frac{1}{y} = \frac{14}{3}$, then $x^3 - y^3$ is equal

to:

- (a) $\frac{433}{64}$ (b) $\frac{217}{32}$
(c) $\frac{217}{64}$ (d) $\frac{433}{32}$

SSC CHSL 19/08/2021 (Shift-II)

Ans. (c) : From,

$$\frac{1}{x} - \frac{1}{y} = \frac{14}{3}$$

$$3(y-x) = 14xy$$

On putting the value of $x-y = \frac{7}{4}$,

$$3\left(\frac{-7}{4}\right) = 14xy$$

$$xy = \frac{-21}{4} \times \frac{1}{14} = \frac{-21}{56}$$

And $x - y = \frac{7}{4}$

On squaring both sides,

$$x^2 + y^2 - 2xy = \frac{49}{16}$$

$$x^2 + y^2 + xy = \frac{49}{16} + 3xy$$

$$x^2 + y^2 + xy = \frac{49}{16} + 3 \times \left(\frac{-21}{56} \right)$$

$$x^2 + y^2 + xy = \frac{49}{16} - \frac{63}{56} = \frac{49 - 18}{16} = \frac{31}{16}$$

Now, $x^3 - y^3 = (x - y)(x^2 + y^2 + xy)$
 $= \frac{7}{4} \times \frac{31}{16} = \frac{217}{64}$

Hence, $x^3 - y^3 = \frac{217}{64}$

122. If $x = 555$, $y = 556$ and $z = 557$, then find the value of $x^3 + y^3 + z^3 - 3xyz$.

- (a) 5006 (b) 5002
(c) 5004 (d) 5008

SSC CHSL 11/08/2021 (Shift-III)

Ans. (c) : From formula

$$x^3 + y^3 + z^3 - 3xyz = \frac{1}{2}(x + y + z)[(x - y)^2 + (y - z)^2 + (z - x)^2]$$

$$x^3 + y^3 + z^3 - 3xyz = \frac{1}{2}(555 + 556 + 557)[(555 - 556)^2 + (556 - 557)^2 + (557 - 555)^2]$$

$$= \frac{1}{2} \times 1668 \times (1 + 1 + 4)$$

$$= \frac{1}{2} \times 1668 \times 6 = 5004$$

Hence $x^3 + y^3 + z^3 - 3xyz = 5004$

123. If $a + 5b = 25$ and $ab = 20$, then one of the values of $(a - 5b)$ is:

- (a) 14 (b) 13
(c) 15 (d) 16

SSC CHSL 11/08/2021 (Shift-III)

Ans. (c) : From formula:-

$$(a + b)^2 - 4ab = (a - b)^2$$

$$(a + 5b)^2 - 4 \times a \times 5b = (a - 5b)^2$$

$$(25)^2 - 20 \times ab = (a - 5b)^2$$

$$625 - 20 \times 20 = (a - 5b)^2$$

$$(a - 5b)^2 = 225$$

$$a - 5b = \sqrt{225} = 15$$

$$a - 5b = 15$$

124. If $3a - b = 1$ and $ab = 4$, then the value of $(9a^2 - b^2)$ is:

- (a) 7 (b) 8
(c) 5 (d) 6

SSC CHSL 11/08/2021 (Shift-III)

Ans. (a) : From formula,

$$(x - y)^2 + 4xy = (x + y)^2$$

$$(3a - b)^2 + 4 \times 3a \times b = (3a + b)^2$$

$$(1)^2 + 12 \times 4 = (3a + b)^2$$

$$3a + b = \sqrt{49} = 7 \text{ ----- (i)}$$

Now, $9a^2 - b^2 = (3a)^2 - (b)^2$

$$9a^2 - b^2 = (3a + b)(3a - b)$$

$$9a^2 - b^2 = 7 \times 1 = 7$$

Hence, $9a^2 - b^2 = 7$

125. If $3x + y = 12$ and $xy = 9$, then the value of $(3x - y)$ is:

- (a) 4 (b) 5
(c) 6 (d) 3

SSC CHSL 09/08/2021 (Shift-III)

Ans. (c) : Given

$$3x + y = 12 \text{ and } xy = 9$$

Now, from $(3x + y)^2 - 4 \times 3x \times y = (3x - y)^2$

$$(12)^2 - 12 \times 9 = (3x - y)^2$$

$$144 - 108 = (3x - y)^2$$

$$36 = (3x - y)^2$$

$$3x - y = \sqrt{36} = 6$$

Hence, $3x - y = 6$

126. If $a^2 + b^2 + c^2 = 576$ and $(ab + bc + ca) = 50$, then what is the value of $(a + b + c)$, if $(a + b + c) < 0$?

- (a) ± 26 (b) -24
(c) -26 (d) ± 24

SSC CHSL 09/08/2021 (Shift-III)

Ans. (c) : From formula,

$$(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$$

$$(a + b + c)^2 = 576 + 2 \times 50$$

[Given $a^2 + b^2 + c^2 = 576$
 $(ab + bc + ca) = 50$]

$$(a + b + c)^2 = 676$$

$$a + b + c = \sqrt{676} = \pm 26$$

$\therefore a + b + c < 0$

Hence, $a + b + c = -26$

127. If $\left(2x + \frac{1}{2x}\right) = 5$, then what is the value of

$$\left(8x^3 + \frac{1}{8x^3}\right)?$$

- (a) 110 (b) 120
(c) 100 (d) 125

SSC CHSL 09/08/2021 (Shift-III)

Ans. (a) : Given $\therefore 2x + \frac{1}{2x} = 5$

On cubing both sides,

$$\left(2x + \frac{1}{2x}\right)^3 = (5)^3$$

$$8x^3 + \frac{1}{8x^3} + 3 \times 2x \times \frac{1}{2x} \left(2x + \frac{1}{2x} \right) = 125$$

$$8x^3 + \frac{1}{8x^3} + 3(5) = 125$$

$$8x^3 + \frac{1}{8x^3} = 125 - 15$$

$$8x^3 + \frac{1}{8x^3} = 110$$

128. If $x + y = 27$ and $x^2 + y^2 = 425$, then the value of $(x - y)^2$ will be:

- (a) 225 (b) 169
(c) 121 (d) 144

SSC CHSL 09/082021 (Shift-II)

Ans. (c) : Given:- $x + y = 27$ and $x^2 + y^2 = 425$

On squaring both side of $x + y = 27$,

$$(x + y)^2 = (27)^2$$

$$x^2 + y^2 + 2xy = 729$$

$$2xy = 729 - 425 = 304 \text{ -----(i)}$$

Now, $(x - y)^2 = x^2 + y^2 - 2xy$
 $= 425 - 304$

$$(x - y)^2 = 121$$

129. If $(40\sqrt{5}x^3 - 2\sqrt{2}y^3) \div (2\sqrt{5}x - \sqrt{2}y) = Ax^2 + By^2 - Cxy$

then find the value of $A + 3B - \sqrt{10}C$

- (a) 34 (b) 46
(c) 6 (d) 28

SSC CHSL 09/082021 (Shift-II)

Ans. (b) :

$$(40\sqrt{5}x^3 - 2\sqrt{2}y^3) \div (2\sqrt{5}x - \sqrt{2}y) = Ax^2 + By^2 - Cxy$$

$$\frac{(2\sqrt{5}x)^3 - (\sqrt{2}y)^3}{2\sqrt{5}x - \sqrt{2}y} = Ax^2 + By^2 - Cxy$$

$$\frac{(2\sqrt{5}x - \sqrt{2}y)(20x^2 + 2y^2 + 2\sqrt{10}xy)}{(2\sqrt{5}x - \sqrt{2}y)} = Ax^2 + By^2 - Cxy$$

$$20x^2 + 2y^2 + 2\sqrt{10}xy = Ax^2 + By^2 - Cxy$$

On comparing both sides,

$$A = 20, B = 2 \text{ and } C = -2\sqrt{10}$$

$$\text{Now } A + 3B - \sqrt{10}C$$

$$= 20 + 3 \times 2 - \sqrt{10} \times (-2\sqrt{10})$$

$$= 20 + 6 + 2 \times 10$$

$$= 26 + 20 = 46$$

130. If $x^4 + \frac{1}{x^4} = 1154, x > 0$, then what will be the value of $x + \frac{1}{x}$?

- (a) $\sqrt{34}$ (b) 18
(c) $\sqrt{32}$ (d) 6

SSC CHSL 09/082021 (Shift-II)

Ans. (d) : Given :-

$$x^4 + \frac{1}{x^4} = 1154$$

$$x^4 + \frac{1}{x^4} + 2 = 1154 + 2$$

$$\left(x^2 + \frac{1}{x^2} \right)^2 = 1156$$

$$\left(x^2 + \frac{1}{x^2} \right) = \sqrt{1156} = 34$$

Now, again adding 2 both sides,

$$x^2 + \frac{1}{x^2} + 2 = 34 + 2$$

$$\left(x + \frac{1}{x} \right)^2 = 36$$

$$x + \frac{1}{x} = \sqrt{36} = 6$$

$$x + \frac{1}{x} = 6$$

131. The value of $a^3 + b^3 + c^3 - 3abc$, when $a = 125, b = 127$ and $c = 129$, is:

- (a) 4725 (b) 4752
(c) 3752 (d) 4572

SSC CHSL 12/08/2021 (Shift-II)

Ans. (d) : From the formula :-

$$a^3 + b^3 + c^3 - 3abc = \frac{(a+b+c)}{2} [(a-b)^2 + (b-c)^2 + (c-a)^2]$$

$$= \left(\frac{125+127+129}{2} \right) [(125-127)^2 + (127-129)^2 + (129-125)^2]$$

$$= 190.5 \times (4+4+16)$$

$$= 190.5 \times 24$$

$$= 4572$$

132. If $(7x + 3)^3 + (x - 2)^3 + 27(2x - 5)^3 = 9(7x + 3)(x - 2)(2x - 5)$, then the value of $5x + 3$ is:

- (a) 2 (b) 10
(c) 6 (d) 8

SSC CHSL 15/04/2021 (Shift-II)

Ans : (d) Given :-

$$(7x + 3)^3 + (x - 2)^3 + 27(2x - 5)^3 = 9(7x + 3)(x - 2)(2x - 5)$$

From formula

If $a^3 + b^3 + c^3 = 3abc$ then $a + b + c$ will be zero

Hence,

$$(7x + 3) + (x - 2) + 3(2x - 5) = 0$$

$$7x + 3 + x - 2 + 6x - 15 = 0$$

$$14x = -3 + 2 + 15 = 14$$

$$x = \frac{14}{14} = 1$$

Now, $5x + 3 = 5 \times 1 + 3 = 8$

$$5x + 3 = 8$$

133. If $(3p - 5m) = 5$ and $pm = 6$, then what is the value of $(9p^2 - 25m^2)$?

- (a) $\pm 30\sqrt{10}$ (b) $30\sqrt{10}$
 (c) $\pm 5\sqrt{385}$ (d) $5\sqrt{385}$

SSC CHSL 15/04/2021 (Shift-II)

Ans. (d) Given:- $(3p - 5m) = 5$, $pm = 6$
 $(3p-5m)^2 + 4 \times 3p \times 5m = (3p+5m)^2$
 $(5)^2 + 60 \times pm = (3p+5m)^2$
 $(3p+5m)^2 = 25 + 360 = 385$

Now, $9p^2 - 25m^2 = (3p)^2 - (5m)^2$
 $= (3p+5m)(3p-5m)$
 $= \sqrt{385} \times 5$

Hence, $9p^2 - 25m^2 = 5\sqrt{385}$

134. If $a + b + c = 2$ and $ab + bc + ca = -1$, then the value of $a^3 + b^3 + c^3 - 3abc$ is:

- (a) 14 (b) 2
 (c) 5 (d) 10

SSC CHSL 06/08/2021 (Shift-II)

Ans. (a) : Given,

$a + b + c = 2 \rightarrow a^2 + b^2 + c^2 + 2(ab+bc+ca) = 4$
 $a^3 + b^3 + c^3 - 3abc = (a+b+c) [a^2 + b^2 + c^2 - ab - bc - ca]$
 $= 2 \times [4 - (ab + bc + ca) \times 2 - (ab+bc+ca)]$
 $= 2 \times [4 - 3(ab+bc+ca)]$
 $= 2 \times [4 + 3]$
 $= 14$

135. If $\left(x^2 + \frac{1}{49x^2}\right) = 15\frac{5}{7}$, then what is the value of

$$\left(x + \frac{1}{7x}\right)?$$

- (a) 7 (b) ± 7
 (c) ± 4 (d) 4

SSC CHSL 06/08/2021 (Shift-II)

Ans. (c) : $x^2 + \frac{1}{49x^2} = 15\frac{5}{7}$

$$\left(x + \frac{1}{7x}\right)^2 = x^2 + \frac{1}{49x^2} + \frac{2}{7} = \frac{110}{7} + \frac{2}{7}$$

$$= \frac{112}{7} = 16$$

$$\therefore \left(x + \frac{1}{7x}\right) = \pm 4$$

136. If $x + \frac{1}{x} = \sqrt{13}$ then findout the value of

$$x^3 - \frac{1}{x^3}.$$

- (a) 32 (b) 36
 (c) $4\sqrt{11}$ (d) $4\sqrt{11}$

SSC CHSL 12/08/2021 (Shift-III)

Ans. (b) : Given

$$x + \frac{1}{x} = \sqrt{13}$$

From formula :-

$$\left(x + \frac{1}{x}\right)^2 - 4 = \left(x - \frac{1}{x}\right)^2$$

$$\left(x - \frac{1}{x}\right)^2 = (\sqrt{13})^2 - 4$$

$$x - \frac{1}{x} = 3$$

On cubing both sides,

$$\left(x - \frac{1}{x}\right)^3 = (3)^3$$

$$x^3 - \frac{1}{x^3} - 3\left(x - \frac{1}{x}\right) = 27$$

$$x^3 - \frac{1}{x^3} - 3 \times 3 = 27$$

$$x^3 - \frac{1}{x^3} = 27 + 9 = 36$$

$$x^3 - \frac{1}{x^3} = 36$$

137. If $x^4 + \frac{16}{x^4} = 27217$, $x > 0$, then the value of $x + \frac{2}{x}$ is:

- (a) 15 (b) 11
 (c) 17 (d) 13

SSC CHSL 15/04/2021 (Shift-III)

Ans.(d) : Given:-

$$x^4 + \frac{16}{x^4} = 27217$$

On adding 8 both sides,

$$\left(x^2\right)^2 + \left(\frac{4}{x^2}\right)^2 + 8 = 27217 + 8$$

$$\left(x^2 + \frac{4}{x^2}\right)^2 = 27225$$

$$x^2 + \frac{4}{x^2} = \sqrt{27225} = 165$$

Again, on adding 4 both sides,

$$\left(x^2\right)^2 + \left(\frac{2}{x}\right)^2 + 4 = 165 + 4 = 169$$

$$\left(x + \frac{2}{x}\right)^2 = 169$$

$$x + \frac{2}{x} = \sqrt{169} = 13$$

Hence $x + \frac{2}{x} = 13$

138. If $8a^3 + b^3 = 16$ and $2a + b = 4$, then find the value of $16a^4 + b^4$.

- (a) 36 (b) 38
(c) 32 (d) 28

SSC CHSL 15/04/2021 (Shift-III)

Ans.(c) : Given:- $8a^3 + b^3 = 16$ -----(i)
and $2a + b = 4$ -----(ii)
From equation (i) and (ii) we will put the value of
 $a = 1$ and $b = 2$
Now $16a^4 + b^4$
 $= 16 \times (1)^4 + (2)^4$
 $= 16 + 16 = 32$
Hence $16a^4 + b^4 = 32$

139. If $x - \frac{1}{2x} = 4$, then the value of $x^3 - \frac{1}{x^3}$ will

- be:**
(a) 480 (b) 540
(c) 520 (d) 560

SSC CHSL 15/04/2021 (Shift-III)

Ans.(d) : Given that- $x - \frac{1}{2x} = 4$
On multiplying by 2 in both sides,
 $2x - \frac{1}{x} = 8$ _____(i)
On cubing both sides,
 $8x^3 - \frac{1}{x^3} - 3 \times 2x \times \frac{1}{x} \left(2x - \frac{1}{x}\right) = 512$
 $\Rightarrow 8x^3 - \frac{1}{x^3} - 6 \times 8 = 512$ [From eqⁿ (i)]
 $\Rightarrow 8x^3 - \frac{1}{x^3} = 512 + 48$
 $8x^3 - \frac{1}{x^3} = 560$

140. If $x^2 + \frac{1}{x^2} = 83$, $x > 0$ then find the value of

- $x^3 - \frac{1}{x^3}$?
(a) 657 (b) 746
(c) 756 (d) 576

SSC CHSL 12/08/2021 (Shift-III)

Ans. (c) : Given, $x^2 + \frac{1}{x^2} = 83$
On subtracting 2 from both sides,
 $x^2 + \frac{1}{x^2} - 2 = 83 - 2 = 81$
 $x - \frac{1}{x} = 9$
On cubing both sides,
 $x^3 - \frac{1}{x^3} = 729 + 27 = 756$

141. If $x^2 + (4 - \sqrt{3})x - 1 = 0$ then find the value of $x^2 + \frac{1}{x^2}$?

- (a) $21 - 12\sqrt{3}$ (b) $17 - 8\sqrt{3}$
(c) $21 - 8\sqrt{3}$ (d) $9 - 8\sqrt{3}$

SSC CHSL 12/08/2021 (Shift-III)

Ans. (c) : Given
 $x^2 + (4 - \sqrt{3})x - 1 = 0$
On multiplying by $\frac{1}{x}$ both sides,
 $x - \frac{1}{x} = 4 - \sqrt{3}$
On squaring both sides,
 $x^2 + \frac{1}{x^2} - 2 = (4 - \sqrt{3})^2 = 16 + 3 - 8\sqrt{3}$
 $x^2 + \frac{1}{x^2} = 19 - 8\sqrt{3} + 2 = 21 - 8\sqrt{3}$
Hence, $x^2 + \frac{1}{x^2} = 21 - 8\sqrt{3}$

142. Given that $(2x+y)^3 - (x+2y)^3 = (x-y)[A(x^2+y^2) + Bxy]$, the value of $(2A - B)$ is:

- (a) 0 (b) 7
(c) 1 (d) 6

SSC CHSL 13/04/2021 (Shift-II)

Ans. (c) : Given:-
 $(2x+y)^3 - (x+2y)^3 = (x-y)[A(x^2+y^2) + Bxy]$
From formula,
 $a^3 - b^3 = (a-b)(a^2 + b^2 + ab)$
 $(2x+y)^3 - (x+2y)^3 = (2x+y - x-2y)[(2x+y)^2 + (x+2y)^2 + (2x+y)(x+2y)]$
 $= (x-y)(5x^2 + 5y^2 + 8xy + 2x^2 + 2y^2 + 5xy)$
 $(x-y)[7(x^2+y^2) + 13xy] = (x-y)[A(x^2+y^2) + Bxy]$
On comparing both sides,
 $A = 7$ and $B = 13$
Now $2A - B = 2 \times 7 - 13$
 $= 14 - 13 = 1$
Hence, $2A - B = 1$

143. If $x^4 - 142x^2 + 1 = 0$, then the value of $x^3 + \frac{1}{x^3}$ is:

- (a) 1592 (b) 1692
(c) 1952 (d) 1962

SSC CHSL 13/04/2021 (Shift-II)

Ans. (b) : Given :-
 $x^4 - 142x^2 + 1 = 0$
 $x^4 + 1 = 142x^2$
 $x^2 + \frac{1}{x^2} = 142$
On adding 2 to both sides -
 $x^2 + \frac{1}{x^2} + 2 = 142 + 2 = 144$

$$\left(x + \frac{1}{x}\right)^2 = (12)^2$$

$$x + \frac{1}{x} = 12$$

On cubing both sides,

$$x^3 + \frac{1}{x^3} + 3\left(x + \frac{1}{x}\right) = 1728$$

$$x^3 + \frac{1}{x^3} + 3 \times 12 = 1728$$

$$x^3 + \frac{1}{x^3} = 1728 - 36$$

$$x^3 + \frac{1}{x^3} = 1692$$

144. If $x^8 - 2599x^4 + 1 = 0$, then the positive value of $x - \frac{1}{x}$ will be:

- (a) 8 (b) 6
(c) 12 (d) 7

SSC CHSL 19/04/2021 (Shift-III)

Ans. (d) : Given, $x^8 - 2599x^4 + 1 = 0$
 $x^8 + 1 = 2599x^4$

On multiplying by $\frac{1}{x^4}$ both sides,

$$x^4 + \frac{1}{x^4} = 2599$$

On adding 2 to both sides,

$$x^4 + \frac{1}{x^4} + 2 = 2599 + 2$$

$$\left(x^2 + \frac{1}{x^2}\right)^2 = (51)^2$$

$$x^2 + \frac{1}{x^2} = 51$$

On subtracting 2 from both sides,

$$x^2 + \frac{1}{x^2} - 2 = 51 - 2 = 49$$

$$\left(x - \frac{1}{x}\right)^2 = (7)^2$$

$$x - \frac{1}{x} = 7$$

145. If $x^2 + \frac{1}{x^2} = 7$, then the value of $x^3 + \frac{1}{x^3}$ where $x > 0$ is equal to :

- (a) 16 (b) 18
(c) 15 (d) 12

SSC CGL (Tier-II)-2019 - 18/11/2020

Ans. (b) : $x^2 + \frac{1}{x^2} = 7$, $x > 0$

$$x^2 + \frac{1}{x^2} + 2 = 9 \quad (\text{adding 2 both side})$$

$$x + \frac{1}{x} = 3$$

$$x^3 + \frac{1}{x^3} = 3^3 - 3 \times 3 = 27 - 9$$

$$\therefore x^3 + \frac{1}{x^3} = 18$$

146. If $x\left(3 - \frac{2}{x}\right) = \frac{3}{x}$ then the value of $x^3 - \frac{1}{x^3}$ is equal to?

- (a) $\frac{8}{27}$ (b) $\frac{52}{27}$
(c) $\frac{62}{27}$ (d) $\frac{61}{27}$

SSC CGL (Tier-II)-2019 - 18/11/2020

Ans. (c) : $x\left(3 - \frac{2}{x}\right) = \frac{3}{x}$

$$3x - 2 = \frac{3}{x}$$

$$x - \frac{1}{x} = \frac{2}{3}$$

From formula if the $x - \frac{1}{x} = K$

$$x^3 - \frac{1}{x^3} = K^3 + 3K$$

Hence,

$$x^3 - \frac{1}{x^3} = \left(\frac{2}{3}\right)^3 + 3 \times \frac{2}{3} \\ = \frac{8}{27} + 2 = \frac{62}{27}$$

147. If $\sqrt{x} + \frac{1}{\sqrt{x}} = 3$, then the value of $x^3 + \frac{1}{x^3}$ is:

- (a) 324 (b) 326
(c) 322 (d) 422

SSC CGL (Tier-II)-2019 - 18/11/2020

Ans. (c) : Let, $\sqrt{x} + \frac{1}{\sqrt{x}} = 3 = K$

$$\therefore x + \frac{1}{x} = K^2 - 2$$

$$x + \frac{1}{x} = 3^2 - 2 = 7$$

$$\therefore x^3 + \frac{1}{x^3} = K^3 - 3K \\ = 7^3 - 3 \times 7 \\ = 343 - 21 = 322$$

148. If $x - \frac{3}{x} = 6$, $x \neq 0$, then the value of

$$\frac{x^4 - \frac{27}{x^2}}{x^2 - 3x - 3}$$

- (a) 90 (b) 80
(c) 270 (d) 54

SSC CGL (Tier-II)-2019 - 18/11/2020

Ans. (a) : $x - \frac{3}{x} = 6, \quad x \neq 0$

$$\left(x - \frac{3}{x}\right)^3 = 6^3 \text{ (On cubing both sides)}$$

$$x^3 - \frac{27}{x^3} - 3 \times x \times \frac{3}{x} \times \left(x - \frac{3}{x}\right) = 216$$

$$x^3 - \frac{27}{x^3} - 9 \times 6 = 216$$

$$x^3 - \frac{27}{x^3} = 270$$

$$\frac{x^4 - \frac{27}{x^2}}{x^2 - 3x - 3} = \frac{x^3 - \frac{27}{x^3}}{x - 3 - \frac{3}{x}}$$

$$= \frac{270}{6 - 3} = 90$$

149. If $\frac{3(x^2+1)-7x}{3x} = 6, x \neq 0$, then the value of $\sqrt{x} + \frac{1}{\sqrt{x}}$ is ?

- (a) $\sqrt{\frac{35}{3}}$ (b) $\sqrt{\frac{31}{3}}$
 (c) $\sqrt{\frac{11}{3}}$ (d) $\sqrt{\frac{25}{3}}$

SSC CGL (Tier-II) 13-09-2019

Ans. (b) :

$$\frac{3(x^2+1)-7x}{3x} = 6$$

$$x + \frac{1}{x} - \frac{7}{3} = 6$$

$$x + \frac{1}{x} = \frac{25}{3}$$

$$x + \frac{1}{x} + 2 = \frac{25}{3} + 2$$

$$\left(\sqrt{x} + \frac{1}{\sqrt{x}}\right)^2 = \frac{31}{3}$$

$$\sqrt{x} + \frac{1}{\sqrt{x}} = \sqrt{\frac{31}{3}}$$

150. If $x+y=3$, then what is the value of x^3+y^3+9xy ?

(a) 15 (b) 81
 (c) 27 (d) 9

SSC CGL (Tier-II) 18-02-2018

Ans. (c) : Given,
 $x + y = 3$
 On cubing both sides
 $x^3 + y^3 + 3xy(x+y) = 27$
 $x^3 + y^3 + 3xy(3) = 27$
 $x^3 + y^3 + 9xy = 27$

Trick:

$$x = 0, y = 3$$

$$x + y = 3$$

$$0 + 3 = 3$$

$$3 = 3 \text{ (equation satisfies)}$$

Now,
 $x^3 + y^3 + 9xy = (0)^3 + (3)^3 + 9 \times 0 \times 3 = 27$

151. If $a^4 + 1 = \left[\frac{a^2}{b^2}\right](4b^2 - b^4 - 1)$, then what is the value of $a^4 + b^4$?

- (a) 2 (b) 16
 (c) 32 (d) 64

SSC CGL (Tier-II) 20-02-2018

Ans. (a)

$$a^4 + 1 = \frac{a^2}{b^2}(4b^2 - b^4 - 1)$$

$$a^2 + \frac{1}{a^2} = 4 - b^2 - \frac{1}{b^2}$$

$$a^2 + \frac{1}{a^2} - 2 + b^2 + \frac{1}{b^2} - 2 = 0$$

$$\left(a - \frac{1}{a}\right)^2 + \left(b - \frac{1}{b}\right)^2 = 0$$

$$\left(a - \frac{1}{a}\right) = 0, \quad \left(b - \frac{1}{b}\right) = 0$$

$$a^2 - 1 = 0, \quad b^2 - 1 = 0$$

$$a = 1, \quad b = 1$$

According to the question

$$a^4 + b^4 = 1 + 1 = 2$$

Trick:

Put, $a = 1, b = 1$

$$a^4 + 1 = \left[\frac{a^2}{b^2}\right][4b^2 - b^4 - 1]$$

$$1 + 1 = 1(4 - 2)$$

$$2 = 2$$

(equation satisfies)

$$\therefore a^4 + b^4 = 1$$

$$\therefore a^4 + b^4 = 1 + 1 = 2$$

152. If $(27x^3 - 343y^3) \div (3x - 7y) = Ax^2 + By^2 + 7Cyx$, then the value of $(4A - B + 5C)$ is:

- (a) 3 (b) 1
 (c) 0 (d) 2

SSC CGL (TIER-I) - 04.06.2019 (Shift-III)

Ans. (d) : $(27x^3 - 343y^3) \div (3x - 7y) = Ax^2 + By^2 + 7Cyx + 7Cyx$

$$a^3 - b^3 = (a-b)(a^2 + b^2 + ab)$$

$$\frac{(3x - 7y)(9x^2 + 49y^2 + 21xy)}{(3x - 7y)} = Ax^2 + By^2 + 7Cyx$$

$$9x^2 + 49y^2 + 21xy = Ax^2 + By^2 + 7Cyx$$

On comparing equation

$$A = 9, B = 49, C = \frac{21}{7} = 3$$

$$4A - B + 5C = 4 \times 9 - 49 + 5 \times 3 = 51 - 49 = 2$$

153. If $a^2 + b^2 + 64c^2 + 16c + 3 = 2(a+b)$, then the value of $4a^7 + b^7 + 8c^2$ is?

- (a) $3\frac{7}{8}$ (b) $5\frac{1}{8}$
 (c) $4\frac{1}{8}$ (d) $4\frac{7}{8}$

SSC CGL (TIER-I)– 04.06.2019 (Shift-III)

Ans. (b) : $a^2 + b^2 + 64c^2 + 16c + 1 + 1 + 1 - 2a - 2b = 0$
 $a^2 - 2a + 1 + b^2 - 2b + 1 + 64c^2 + 16c + 1 = 0$
 $(a-1)^2 + (b-1)^2 + (8c+1)^2 = 0$

$(a-1)^2 = 0$	$(b-1)^2 = 0$	$(8c+1)^2 = 0$
$a = 1$	$b = 1$	$8c = -1$
		$c = -\frac{1}{8}$

$\Rightarrow 4a^7 + b^7 + 8c^2 = 4 \times 1 + 1 + 8 \times \frac{1}{64}$
 $= 4 + 1 + \frac{1}{8} = 5\frac{1}{8}$

154. If $x + y = 1$ and $xy(xy - 2) = 12$, then the value of $x^4 + y^4$ is:

- (a) 19 (b) 23
 (c) 25 (d) 20

SSC CGL (TIER-I) – 04.06.2019 (Shift-III)

Ans. (c) : $x + y = 1$
 On squaring both sides
 $x^2 + y^2 + 2xy = 1$
 $x^2 + y^2 = 1 - 2xy$
 Again, on squaring both sides
 $x^4 + y^4 + 2x^2y^2 = 1 + 4x^2y^2 - 4xy$
 $x^4 + y^4 = 1 + 2xy(xy - 2)$ [$\because xy(xy - 2) = 12$]
 $= 1 + 2 \times 12$
 $= 25$

155. If $a^2 + b^2 + c^2 = 21$, and $a + b + c = 7$, then $(ab + bc + ca)$ is equal to?

- (a) 14 (b) 8
 (c) 12 (d) 28

SSC CGL (TIER-I) – 04.06.2019 (Shift-III)

Ans. (a) : $(a + b + c)^2 = a^2 + b^2 + c^2 + 2(ab + bc + ca)$
 $49 = 21 + 2(ab + bc + ca)$
 $\frac{28}{2} = (ab + bc + ca)$
 $(ab + bc + ca) = 14$

156. If $16x^2 + 9y^2 + 4z^2 = 24(x - y + z) - 61$, then the value of $(xy + 2z)$ is:

- (a) 1 (b) 2
 (c) 5 (d) 3

SSC CGL (TIER-I) – 04.06.2019 (Shift-II)

Ans. (c) : Given that,
 $16x^2 + 9y^2 + 4z^2 = 24(x - y + z) - 61$
 $[xy + 2z = ?]$
 $16x^2 + 9y^2 + 4z^2 - 24x + 24y - 24z + 61 = 0$
 $(16x^2 - 24x + 9) + (9y^2 + 24y + 16) + (4z^2 - 24z + 36) = 0$

$$(4x - 3)^2 + (3y + 4)^2 + (2z - 6)^2 = 0$$

$$4x - 3 = 0 \quad | \quad 3y + 4 = 0 \quad | \quad 2z - 6 = 0$$

$$x = 3/4 \quad | \quad y = -4/3 \quad | \quad z = 3$$

$$\Rightarrow xy + 2z = \frac{3}{4} \times \left(-\frac{4}{3}\right) + 2 \times 3$$

$$= -1 + 6 = 5$$

157. If $[8(x + y)^3 - 27(x - y)^3] \div (5y - x) = Ax^2 + Bxy + Cy^2$, then the value of $(A + B + C)$ is?

- (a) 26 (b) 19
 (c) 13 (d) 16

SSC CGL (TIER-I)– 04.06.2019 (Shift-II)

Ans. (d) :
 $[8(x + y)^3 - 27(x - y)^3] \div (5y - x) = Ax^2 + Bxy + Cy^2$
 $[A + B + C = ?]$
 $a^3 - b^3 = (a - b)(a^2 + b^2 + ab)$
 $[2(x + y)]^3 - [3(x - y)]^3 \div (5y - x) = Ax^2 + Bxy + Cy^2$
 $\Rightarrow \frac{(5y - x)[2(x + y)]^2 + [3(x - y)]^2 + 2(x + y) \times (3(x - y))}{(5y - x)}$
 $= Ax^2 + Bxy + Cy^2$
 $\Rightarrow \frac{(5y - x)[2(x + y)]^2 + [3(x - y)]^2 + 6(x^2 - y^2)}{(5y - x)}$
 $= Ax^2 + Bxy + Cy^2$
 $= 4(x + y)^2 + 9(x - y)^2 + 6(x^2 - y^2) = Ax^2 + Bxy + Cy^2$
 $4(x^2 + y^2) + 9(x^2 + y^2) + 6(x^2 - y^2) + 8xy - 18xy = Ax^2 + Bxy + Cy^2$
 $19x^2 + 7y^2 - 10xy = Ax^2 + Bxy + Cy^2$
 On comparing both sides with the respective term
 $A = 19$
 $B = -10$
 $C = 7$
 $\Rightarrow A + B + C = 19 + 7 - 10 = 16$

158. If $x + y + z = 19$, $xy + yz + zx = 114$, then the value of $\sqrt{x^3 + y^3 + z^3 - 3xyz}$ is:

- (a) 17 (b) 13
 (c) 19 (d) 21

SSC CGL (TIER-I) – 04.06.2019 (Shift-II)

Ans. (c) $(x + y + z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)$
 $361 = x^2 + y^2 + z^2 + 2 \times 114$
 $361 - 228 = x^2 + y^2 + z^2$
 $x^2 + y^2 + z^2 = 133$
 $x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)$
 $= 19 \times (133 - 114)$
 $\sqrt{x^3 + y^3 + z^3 - 3xyz} = \sqrt{19 \times 19}$
 $\sqrt{x^3 + y^3 + z^3 - 3xyz} = 19$

159. If $x + y + z = 19$, $x^2 + y^2 + z^2 = 133$ and $xz = y^2$, then the difference between z and x is:

- (a) 6 (b) 5
 (c) 3 (d) 4

SSC CGL (TIER-I)– 04.06.2019 (Shift-I)

Ans. (b) $x + y + z = 19$ (i)
 $x^2 + y^2 + z^2 = 133$ (ii)
 $xz = y^2$ (iii)
 $z - x = ?$
 $(x+y+z) = 19$ (On squaring both sides)
 $x^2 + y^2 + z^2 + 2xy + 2yz + 2xz = 361$
 $133 + 2(xy + yz + zx) = 361$
 $2(xy + yz + zx) = 361 - 133$
 $xy + yz + zx = 114$
 $xz = y^2$ (On putting the value) from equation (iii)
 $xy + yz + y^2 = 114$
 $y(x+y+z) = 114$ ($x+y+z = 19$ From equation (i))
 $y \times 19 = 114$
 $y = 6$
 $y^2 = xz$
 $36 = x \times z$
 $36 = 4 \times 9$
 $36 = 36$
Hence, we will put the value of x and z in such a way that the equation $x + y + z = 19$ is satisfied.
 $x = 4$
 $y = 6$
 $z = 9$
 $\therefore z - x = 9 - 4 = 5$

OR

$x + y + z = 19$, $x^2 + y^2 + z^2 = 133$, $xz = y^2$, $z - x = ?$
Equation is satisfied from the value of $x = 4$, $y = 6$, $z = 9$
 $x + y + z = 19$
 $4 + 6 + 9 = 19$
 $19 = 19$
 $\therefore z - x = 9 - 4 = 5$

- 160. If** $3\sqrt{3}x^3 - 2\sqrt{2}y^3 = (\sqrt{3}x - \sqrt{2}y)(Ax^2 + By^2 + Cxy)$,
then the value of $(A \times B) \div C$ is?
(a) $\sqrt{3}$ (b) $\sqrt{6}$
(c) $6\sqrt{6}$ (d) $6\sqrt{3}$
SSC CGL (TIER-I) - 06.06.2019 (Shift-III)

Ans. (b) :
 $3\sqrt{3}x^3 - 2\sqrt{2}y^3 = (\sqrt{3}x - \sqrt{2}y)(Ax^2 + By^2 + Cxy)$
 $(\sqrt{3}x)^3 - (\sqrt{2}y)^3 = (\sqrt{3}x - \sqrt{2}y)(Ax^2 + By^2 + Cxy)$
 $(\sqrt{3}x - \sqrt{2}y)(3x^2 + 2y^2 + \sqrt{6}xy) = (\sqrt{3}x - \sqrt{2}y)(Ax^2 + By^2 + Cxy)$
 $3x^2 + 2y^2 + \sqrt{6}xy = Ax^2 + By^2 + Cxy$
By comparing
 $A = 3$, $B = 2$, $C = \sqrt{6}$
Hence $(A \times B) \div C = (3 \times 2) \div \sqrt{6} = \sqrt{6}$

- 161. If $a + b + c = 2$, $a^2 + b^2 + c^2 = 26$, then the value of $a^3 + b^3 + c^3 - 3abc$ is?**
(a) 71 (b) 74
(c) 78 (d) 69
SSC CGL (TIER-I) - 06.06.2019 (Shift-III)

Ans. (b) : $\because (a + b + c)^2 = a^2 + b^2 + c^2 + 2(ab + bc + ca)$
 $4 = 26 + 2(ab + bc + ca)$
 $ab + bc + ca = -11$
 $a^3 + b^3 + c^3 - 3abc = (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ca)$
 $= 2(26 + 11)$
 $= 2 \times 37 = 74$

- 162. If $a + \frac{1}{a} = 3$, then $\left(a^4 + \frac{1}{a^4}\right)$ is equal to:**
(a) 47 (b) 27
(c) 77 (d) 81
SSC CGL (TIER-I) - 06.06.2019 (Shift-III)

Ans. (a) : $a + \frac{1}{a} = 3$
 $\left(a + \frac{1}{a}\right)^2 = 9$ (On squaring both sides)
 $a^2 + \frac{1}{a^2} + 2 = 9$
 $a^2 + \frac{1}{a^2} = 7$ (Again, on squaring both side)
 $\left(a^2 + \frac{1}{a^2}\right)^2 = 49$
 $a^4 + \frac{1}{a^4} + 2 = 49$
 $a^4 + \frac{1}{a^4} = 47$

- 163. If $x = a + \frac{1}{a}$ and $y = a - \frac{1}{a}$ then $\sqrt{x^4 + y^4 - 2x^2y^2}$ is equal to?**
(a) $16a^2$ (b) $\frac{8}{a^2}$
(c) 4 (d) 8
SSC CGL (TIER-I) - 06.06.2019 (Shift-I)

Ans. (c) : $\sqrt{x^4 + y^4 - 2x^2y^2}$
 $= \sqrt{(x^2 - y^2)^2}$
 $= x^2 - y^2$
 $= (x + y)(x - y)$
 $= 2a \times \frac{2}{a} = 4$

- 164. If $ab + bc + ca = 8$ and $a^2 + b^2 + c^2 = 20$, then a possible value of $\frac{1}{2}(a + b + c) \left[(a - b)^2 + (b - c)^2 + (c - a)^2 \right]$ is:**
(a) 84 (b) 56
(c) 72 (d) 80
SSC CGL (TIER-I) - 06.06.2019 (Shift-I)

Ans. (c) : $\because (a+b+c)^2 = a^2 + b^2 + c^2 + 2(ab+bc+ca)$
 $= 20 + 2 \times 8 = 36$
 $a+b+c = 6$
 $\therefore \frac{1}{2}(a+b+c)[(a-b)^2 + (b-c)^2 + (c-a)^2]$
 $= (a+b+c)(a^2 + b^2 + c^2 - ab - bc - ca)$
 $= 6 \times [20 - 8]$
 $= 6 \times 12 = 72$

165. If $\frac{6x}{(2x^2 + 5x - 2)} = 1, x > 0$, then the value of $x^3 + \frac{1}{x^3}$ is?
- (a) $\frac{3}{8}\sqrt{17}$ (b) $\frac{5\sqrt{17}}{8}$
(c) $\frac{5\sqrt{17}}{16}$ (d) $\frac{3}{4}\sqrt{17}$

SSC CGL (TIER-I)-07.06.2019 (Shift-III)

Ans. (b) : Given, $\frac{6x}{(2x^2 + 5x - 2)} = 1$
 $\Rightarrow \frac{6}{2x + 5 - \frac{2}{x}} = 1 \Rightarrow 2\left(x - \frac{1}{x}\right) = 1$
 $\Rightarrow \left(x - \frac{1}{x}\right) = \frac{1}{2}$
 $\therefore \left(x + \frac{1}{x}\right) = \sqrt{\left(x - \frac{1}{x}\right)^2 + 4}$
 $= \sqrt{\left(\frac{1}{2}\right)^2 + 4} = \frac{\sqrt{17}}{2}$
 $\therefore x^3 + \frac{1}{x^3} = \left(x + \frac{1}{x}\right)\left[x^2 + \frac{1}{x^2} - 1\right]$
 $= \left(x + \frac{1}{x}\right)\left[\left(x + \frac{1}{x}\right)^2 - 2\right] - 1$
 $= \left(\frac{\sqrt{17}}{2}\right)\left[\left(\frac{\sqrt{17}}{2}\right)^2 - 3\right]$
 $= \frac{\sqrt{17}}{2}\left[\frac{17-12}{4}\right]$
 $= \frac{5\sqrt{17}}{8}$

166. If $x^2 - 3x - 1 = 0$, then the value of $(x^2 + 8x - 1)(x^3 + x^{-1})^{-1}$ is?

- (a) 8 (b) $\frac{3}{8}$
(c) 1 (d) 3

SSC CGL (TIER-I)-2018 - 07.06.2019 (Shift-II)

Ans. (c) : $x^2 - 3x - 1 = 0$
 $x^2 - 1 = 3x$ (i)
 $(x^2 - 1)^2 = (3x)^2$ (On squaring both sides)
 $x^4 + 1 - 2x^2 = 9x^2$
 $x^4 + 1 = 11x^2$ (ii)
 $(x^2 + 8x - 1)\left(\frac{x^4 + 1}{x}\right)^{-1}$

From question,

$(11x)\left(\frac{11x^2}{x}\right)^{-1} \quad [\because x^2 - 1 = 3x]$
 $(11x) \times \left(\frac{1}{11x}\right) = 1$

167. If $(135\sqrt{5}x^3 - 2\sqrt{2}y^3) \div (3\sqrt{5}x - \sqrt{2}y) = Ax^2 + By^2 + \sqrt{10}Cxy$, then the value of $(A + B - 9C)$ is?

- (a) 18 (b) 12
(c) 10 (d) 20

SSC CGL (TIER-I)-2018 - 07.06.2019 (Shift-II)

Ans. (d) : $(135\sqrt{5}x^3 - 2\sqrt{2}y^3) \div (3\sqrt{5}x - \sqrt{2}y)$
 $= Ax^2 + By^2 + \sqrt{10}Cxy$
 $\left[(3\sqrt{5}x)^3 - (\sqrt{2}y)^3\right] \div (3\sqrt{5}x - \sqrt{2}y)$
 $= Ax^2 + By^2 + \sqrt{10}Cxy$
 $\therefore a^3 - b^3 = (a-b)(a^2 + b^2 + ab)$
 $(3\sqrt{5}x - \sqrt{2}y)(45x^2 + 2y^2 + 3\sqrt{10}xy) \div (3\sqrt{5}x - \sqrt{2}y)$
 $= Ax^2 + By^2 + \sqrt{10}Cxy$
 $45x^2 + 2y^2 + 3\sqrt{10}xy = Ax^2 + By^2 + \sqrt{10}Cxy$
By comparing
 $A = 45, B = 2, C = 3$
 $(A + B - 9C) = 45 + 2 - 27$
 $= 47 - 27$
 $= 20$

168. If $9a^2 + 4b^2 + c^2 + 21 = 4(3a + b - 2c)$, then the value of $(9a + 4b - c)$ is?

- (a) 12 (b) 2
(c) 16 (d) 6

SSC CGL (TIER-I)-2018 - 07.06.2019 (Shift-II)

Ans. (a) : $9a^2 + 4b^2 + c^2 + 21 = 4(3a + b - 2c)$
 $(9a^2 - 12a + 4) + (4b^2 - 4b + 1) + (c^2 + 8c + 16) = 0$
 $(3a - 2)^2 + (2b - 1)^2 + (c + 4)^2 = 0$
 $3a - 2 = 0 \quad 2b - 1 = 0 \quad c + 4 = 0$

$$\Rightarrow a = 2/3 \Rightarrow b = 1/2 \Rightarrow c = -4$$

$$9a + 4b - c$$

$$= 9 \times \frac{2}{3} + 4 \times \frac{1}{2} - (-4)$$

$$= 6 + 2 + 4 = 12$$

169. If $x = 2 - p$, then $x^3 + 6xp + p^3$ is equal to?

- (a) 12 (b) 6
(c) 8 (d) 4

SSC CGL (TIER-I)-2018 – 07.06.2019 (Shift-I)

Ans. (c) : $x = 2 - p$

$$x + p = 2$$

On cubing both sides

$$(x + p)^3 = (2)^3$$

$$x^3 + p^3 + 3xp(x + p) = 8$$

$$x^3 + p^3 + 3xp(2) = 8$$

$$\boxed{x^3 + p^3 + 6xp = 8}$$

170. If $x^4 - 6x^2 - 1 = 0$, then the value of

$$x^6 - 5x^2 + \frac{5}{x^2} - \frac{1}{x^6} + 5 \text{ is?}$$

- (a) 239 (b) 204
(c) 209 (d) 219

SSC CGL (TIER-I)-2018 – 07.06.2019 (Shift-I)

Ans. (c) :

$$x^6 - 5x^2 + \frac{5}{x^2} - \frac{1}{x^6} + 5$$

$$x^6 - \frac{1}{x^6} - 5\left(x^2 - \frac{1}{x^2}\right) + 5 \quad \dots\dots\dots(A)$$

$$\therefore x^4 - 6x^2 - 1 = 0$$

$$x^4 - 1 = 6x^2$$

$$x^2 - \frac{1}{x^2} = 6 \quad \dots\dots\dots(B)$$

$$\left(x^2 - \frac{1}{x^2}\right)^3 = (6)^3$$

$$x^6 - \frac{1}{x^6} - 3\left(x^2 - \frac{1}{x^2}\right) = 216$$

$$x^6 - \frac{1}{x^6} - 3(6) = 216$$

$$x^6 - \frac{1}{x^6} = 216 + 18$$

$$\boxed{x^6 - \frac{1}{x^6} = 234} \quad \dots\dots\dots(C)$$

By putting the value of equation B and equation (C) in equation (A)

$$= 234 - 5(6) + 5$$

$$= 234 - 30 + 5$$

$$= \boxed{209}$$

171. If $a + b + c = 11$ and $ab + bc + ca = 38$, then $a^3 + b^3 + c^3 - 3abc$ is equal to?

- (a) 44 (b) 77
(c) 55 (d) 66

SSC CGL (TIER-I)-2018 – 10.06.2019 (Shift-III)

Ans. (b) : $a + b + c = 11 \dots\dots\dots(i)$

Given

$$\therefore ab + bc + ca = 38$$

From equation (i)

$$(a + b + c)^2 = (11)^2 = 121$$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = 121$$

$$a^2 + b^2 + c^2 = 121 - 2 \times 38 = 121 - 76 = 45$$

$$\therefore a^3 + b^3 + c^3 - 3abc = (a+b+c)(a^2+b^2+c^2-ab-bc-ca)$$

$$\therefore a^3 + b^3 + c^3 - 3abc = 11 \times (45 - 38)$$

$$= 11 \times 7$$

$$= 77$$

172. If $\sqrt{x} - \frac{1}{\sqrt{x}} = 4$, then $x^2 + \frac{1}{x^2}$ is equal to?

- (a) 192 (b) 322
(c) 256 (d) 326

SSC CGL (TIER-I)-2018 – 10.06.2019 (Shift-II)

Ans. (b) : Given,

$$\sqrt{x} - \frac{1}{\sqrt{x}} = 4$$

On squaring both sides

$$\left(\sqrt{x} - \frac{1}{\sqrt{x}}\right)^2 = (4)^2$$

$$x + \frac{1}{x} - 2 = 16$$

$$\left(x + \frac{1}{x}\right) = 18 \quad \dots\dots\dots(i)$$

$$\therefore x^2 + \frac{1}{x^2} = \left(x + \frac{1}{x}\right)^2 - 2$$

$$x^2 + \frac{1}{x^2} = (18)^2 - 2 \quad \text{(From equation (i))}$$

$$x^2 + \frac{1}{x^2} = 324 - 2$$

$$x^2 + \frac{1}{x^2} = 322$$

173. If $4x^2 - 6x + 1 = 0$, then the value of $8x^3 + (8x^3)^{-1}$ is:

- (a) 36 (b) 13
(c) 11 (d) 18

SSC CGL (TIER-I)-2018 – 10.06.2019 (Shift-I)

Ans. (d) : $4x^2 - 6x + 1 = 0$

$$4x^2 + 1 = 6x$$

$$\frac{4x^2 + 1}{2x} = \frac{6x}{2x} \quad (\because \text{on dividing by } 2x)$$

$$2x + \frac{1}{2x} = 3 \quad \dots\dots\dots(i)$$

From equation (i)

$$\left(2x + \frac{1}{2x}\right)^3 = 8x^3 + \frac{1}{8x^3} + 3 \times 2x \times \frac{1}{2x} \left(2x + \frac{1}{2x}\right)$$

$$(3)^3 = 8x^3 + \frac{1}{8x^3} + 9$$

$$18 = 8x^3 + \frac{1}{8x^3}$$

or, $8x^3 + (8x^3)^{-1} = 18$

174. If $\sqrt{x} + \frac{1}{\sqrt{x}} = \sqrt{7}$, then $x^3 + \frac{1}{x^3}$ is equal to?

- (a) 120 (b) 110
(c) 140 (d) 130

SSC CGL (TIER-I)-2018 – 11.06.2019 (Shift-III)

Ans. (b) $\sqrt{x} + \frac{1}{\sqrt{x}} = \sqrt{7}$

$$x + \frac{1}{x} + 2 = 7 \quad (\text{By squaring both sides})$$

$$x + \frac{1}{x} = 5$$

$$[(a+b)^3 = a^3 + b^3 + 3ab(a+b)]$$

$$\left(x + \frac{1}{x}\right)^3 = (5)^3$$

$$x^3 + \frac{1}{x^3} + 3 \times x \times \frac{1}{x} \left(x + \frac{1}{x}\right) = 125$$

$$x^3 + \frac{1}{x^3} + 3(5) = 125, \quad \boxed{x^3 + \frac{1}{x^3} = 110}$$

175. If $a + b = 8$ and $ab = \frac{32}{3}$, then $(a^3 + b^3)$ is equal to:

- (a) 256 (b) 384
(c) 128 (d) 320

SSC CGL (TIER-I)-2018 – 11.06.2019 (Shift-III)

Ans. (a) : We know that formula

$$[(a+b)^3 = a^3 + b^3 + 3ab(a+b)]$$

According to this formula

$$(8)^3 = a^3 + b^3 + 3 \times \frac{32}{3} (8)$$

$$512 = a^3 + b^3 + 256$$

$$a^3 + b^3 = 256$$

176. If $a + b + c = 4$ and $ab + bc + ca = 2$, then $a^3 + b^3 + c^3 - 3abc$ is equal to:

- (a) 36 (b) 32
(c) 48 (d) 40

SSC CGL (TIER-I)-2018 – 11.06.2019 (Shift-III)

Ans. (d) : We know that:-

$$[(a+b+c)^2 = a^2 + b^2 + c^2 + 2(ab+bc+ca)]$$

$$(4)^2 = a^2 + b^2 + c^2 + 2(2)$$

$$a^2 + b^2 + c^2 = 12$$

$$a^3 + b^3 + c^3 - 3abc = (a+b+c) [a^2 + b^2 + c^2 - (ab + bc + ca)] = 4 [12 - 2]$$

$$a^3 + b^3 + c^3 - 3abc = 4 \times 10 = 40$$

177. If $(a + b) = 6$ and $ab = \frac{16}{3}$, then $(a^3 + b^3)$ is

equal to:

- (a) 150 (b) 190
(c) 220 (d) 120

SSC CGL (TIER-I)-2018 – 11.06.2019 (Shift-II)

Ans. (d) : Given,

$$a + b = 6 \quad \dots(i)$$

$$ab = \frac{16}{3}$$

$$\therefore a^3 + b^3 = (a + b)(a^2 + b^2 - ab) \quad \dots(ii)$$

From equation (i)

$$(a + b)^2 = 36$$

$$a^2 + b^2 + 2ab = 36$$

$$a^2 + b^2 = 36 - 2 \times \frac{16}{3} = \frac{108 - 32}{3} = \frac{76}{3}$$

From equation (ii)

$$a^3 + b^3 = 6 \times \left(\frac{76}{3} - \frac{16}{3}\right) = 6 \times \frac{60}{3} = 120$$

178. If $a + b + c = 8$ and $ab + bc + ca = 12$, then $a^3 + b^3 + c^3 - 3abc$ is equal to:

- (a) 192 (b) 144
(c) 400 (d) 224

SSC CGL (TIER-I)-2018 – 11.06.2019 (Shift-I)

Ans. (d) : $a + b + c = 8$, $ab + bc + ca = 12$ Given that

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = 64$$

$$a^2 + b^2 + c^2 = 64 - 2(ab + bc + ca) \quad \dots(i)$$

Now,

$$a^3 + b^3 + c^3 - 3abc = (a+b+c)(a^2 + b^2 + c^2 - ab - bc - ca)$$

\therefore From equation (i)

$$a^3 + b^3 + c^3 - 3abc = (a+b+c) [64 - 3(ab+bc+ca)]$$

$$= 8 \times [64 - 3 \times 12] = 8 \times [64 - 36]$$

$$= 8 \times 28$$

$$= 224$$

179. If $a - b = 5$ and $ab = 2$, then $a^3 - b^3$ is equal to?

- (a) 95 (b) 145
(c) 125 (d) 155

SSC CGL (TIER-I)-2018 – 12.06.2019 (Shift-III)

Ans. (d) : $a - b = 5$, $ab = 2$

$$a^3 - b^3 = (a - b)^3 + 3ab(a - b)$$

$$a^3 - b^3 = (5)^3 + 3 \times 2 \times (5)$$

$$= 125 + 30$$

$$= 155$$

180. If $\sqrt{x} - \frac{1}{\sqrt{x}} = 2\sqrt{2}$, then $x^2 + \frac{1}{x^2}$ is equal to:

- (a) 100 (b) 98
(c) 102 (d) 104

SSC CGL (TIER-I)-2018 – 12.06.2019 (Shift-II)

Ans. (b) :

$$\sqrt{x} - \frac{1}{\sqrt{x}} = 2\sqrt{2} \quad \dots(i)$$

On squaring equation (i)

$$\left(\sqrt{x} - \frac{1}{\sqrt{x}}\right)^2 = (2\sqrt{2})^2$$

$$x + \frac{1}{x} - 2 = 8$$

$$x + \frac{1}{x} = 10 \quad \dots\dots(ii)$$

On squaring equation (ii)

$$\left(x + \frac{1}{x}\right)^2 = (10)^2$$

$$x^2 + \frac{1}{x^2} = 98$$

181. If $(a + b) = 6$ and $ab = 8$, then $(a^3 + b^3)$ is equal to?

- (a) 72 (b) 108
(c) 144 (d) 216

SSC CGL (TIER-I)-2018 – 12.06.2019 (Shift-I)

Ans. (a) :

$$(a + b) = 6$$

by cubing both sides

$$(a+b)^3 = (6)^3$$

$$\Rightarrow a^3 + b^3 + 3ab(a+b) = 216$$

$$\Rightarrow a^3 + b^3 + 3 \times 8 \times 6 = 216 \quad [\because ab = 8]$$

$$\Rightarrow a^3 + b^3 = 216 - 144$$

$$\Rightarrow \boxed{a^3 + b^3 = 72}$$

182. If $x + \frac{1}{x} = 5$, then $x^3 + \frac{1}{x^3}$ is equal to:

- (a) 110 (b) 130
(c) 125 (d) 145

SSC CGL (TIER-I)-2018 – 13.06.2019 (Shift-III)

Ans. (a) : Given-

$$x + \frac{1}{x} = 5$$

$$\therefore x^3 + \frac{1}{x^3} = \left(x + \frac{1}{x}\right)^3 - 3\left(x + \frac{1}{x}\right) \\ = 125 - 15 = 110$$

183. If $(x-5)^3 + (x-6)^3 + (x-7)^3 = 3(x-5)(x-6)(x-7)$ then what is the value of x ?

- (a) 18 (b) 6
(c) 5 (d) 7

SSC CGL (TIER-I)-2018 – 13.06.2019 (Shift-III)

$$\text{Ans. (b) : } (x-5)^3 + (x-6)^3 + (x-7)^3 \\ = 3(x-5)(x-6)(x-7)$$

$$a = x - 5, b = x - 6, c = x - 7$$

$$\therefore \boxed{a^3 + b^3 + c^3 = 3abc}$$

$$\therefore a + b + c = 0$$

$$x - 5 + x - 6 + x - 7 = 0$$

$$3x - 18 = 0$$

$$x = 6$$

184. If $a^3 - b^3 = 208$ and $a - b = 4$, then $(a + b)^2 - ab$ is equal to:

- (a) 32 (b) 38
(c) 52 (d) 42

SSC CGL (TIER-I)-2018 – 13.06.2019 (Shift-III)

$$\text{Ans. (c) : } a^3 - b^3 = 208$$

$$(a - b)(a^2 + b^2 + ab) = 208$$

$$4(a^2 + b^2 + ab) = 208$$

$$a^2 + b^2 + ab = 52$$

$$\therefore (a + b)^2 - ab = 52$$

185. If $a - b = 5$ and $ab = 6$, then $(a^3 - b^3)$ is equal to?

- (a) 215 (b) 155
(c) 90 (d) 225

SSC CGL (TIER-I)-2018 – 13.06.2019 (Shift-II)

Ans. (a) : $a - b = 5$ and $ab = 6$

$$\boxed{a^3 - b^3 = (a - b)^3 + 3ab(a - b)}$$

$$a^3 - b^3 = 5^3 + 3 \times 6 \times 5$$

$$a^3 - b^3 = 125 + 90$$

$$a^3 - b^3 = 215$$

186. If $a - \frac{1}{a} = 3$, then $a^6 + \frac{1}{a^6}$ equal to :

- (a) 996 (b) 729
(c) 1298 (d) 1331

SSC CGL (TIER-I)-2018 – 19.06.2019 (Shift-III)

$$\text{Ans. (c) : } a - \frac{1}{a} = 3 \dots\dots(i)$$

By squaring the equation (i)

$$a^2 + \frac{1}{a^2} - 2 = 9$$

$$a^2 + \frac{1}{a^2} = 11 \dots\dots(ii)$$

By cubing the equation (ii)

$$a^6 + \frac{1}{a^6} + 3\left(a^2 + \frac{1}{a^2}\right) = 1331$$

$$a^6 + \frac{1}{a^6} = 1331 - 33 = 1298$$

187. If $x + \frac{1}{x} = 4$, then $x^3 + \frac{1}{x^3}$ equal to :

- (a) 52 (b) 64
(c) 40 (d) 50

SSC CGL (TIER-I)-2018 – 19.06.2019 (Shift-III)

$$\text{Ans. (a) : } x + \frac{1}{x} = 4$$

$$\boxed{x^3 + \frac{1}{x^3} = a^3 - 3a}$$

On cubing both sides

$$x^3 + \frac{1}{x^3} = (4)^3 - 3 \times 4 = 64 - 12$$

$$\therefore x^3 + \frac{1}{x^3} = 52$$

188. If $a^3 - b^3 = 210$ and $a - b = 5$, then $(a+b)^2 - ab$ equal to:

- (a) 52 (b) 42
(c) 38 (d) 32

SSC CGL (TIER-I)-2018 – 19.06.2019 (Shift-III)

$$\text{Ans. (b) } \therefore a^3 - b^3 = 210, a - b = 5$$

$$(a - b)(a^2 + b^2 + ab) = 210$$

$$\therefore a - b = 5 \text{ Given}$$

$$\therefore 5 \times (a^2 + b^2 + ab) = 210$$

$$(a^2 + b^2 + 2ab) - ab = 42$$

$$(a + b)^2 - ab = 42$$

189. If $(x-4)^3 + (x-5)^3 + (x-3)^3 = 3(x-4)(x-5)(x-3)$, then what will be the value of x.

- (a) 7 (b) 4
(c) 18 (d) 6

SSC CGL (TIER-I)-2018 – 19.06.2019 (Shift-III)

Ans. (b) $a^3 + b^3 + c^3 = 3abc$ It is possible only when $a + b + c = 0$
 $x-4 + x-5 + x-3 = 0$
 $3x - 12 = 0$
 $3x = 12 \quad x = 4$

IInd method

$$(x-4)^3 + (x-5)^3 + (x-3)^3 = 3(x-4)(x-5)(x-3)$$

∴ By option (b)

∴ x = 4 By taking the value of x = 4

$$(4-4)^3 + (4-5)^3 + (4-3)^3 = 3(4-4)(4-5)(4-3)$$

$$0 + (-1) + 1 = 3(0) \times (-1)(1)$$

$$0 = 0$$

$$\text{L.H.S.} = \text{R.H.S.}$$

$$\therefore x = 4$$

190. If $x^2 + 3x + 1 = 0$, then what is the value of

$$x^6 + \frac{1}{x^6} ?$$

- (a) 324 (b) 327
(c) 322 (d) 318

SSC CGL (Tier-I)-2019 – 03/03/2020 (Shift-II)

Ans. (c) : $x^2 + 3x + 1 = 0$

Dividing by x to both sides

$$x + 3 + \frac{1}{x} = 0$$

$$x + \frac{1}{x} = -3$$

On cubing both sides,

$$\left(x + \frac{1}{x}\right)^3 = (-3)^3$$

$$x^3 + \frac{1}{x^3} + 3(x) \frac{1}{(x)} \left(x + \frac{1}{x}\right) = -27$$

$$x^3 + \frac{1}{x^3} + 3(-3) = -27$$

$$x^3 + \frac{1}{x^3} = -27 + 9$$

$$x^3 + \frac{1}{x^3} = -18$$

On squaring both sides

$$\left(x^3 + \frac{1}{x^3}\right)^2 = (-18)^2$$

$$x^6 + \frac{1}{x^6} + 2 = 324$$

$$x^6 + \frac{1}{x^6} = 324 - 2 = 322$$

191. The value of $27a^3 - 2\sqrt{2}b^3$ is equal to?

- (a) $(3a - \sqrt{2}b)(9a^2 + 2b^2 + 6\sqrt{2}ab)$
(b) $(3a - \sqrt{2}b)(9a^2 - 2b^2 - 3\sqrt{2}ab)$

(c) $(3a - \sqrt{2}b)(9a^2 + 2b^2 + 3\sqrt{2}ab)$

(d) $(3a - \sqrt{2}b)(9a^2 - 2b^2 + 6\sqrt{2}ab)$

SSC CGL (Tier-I)-2019 – 03/03/2020 (Shift-III)

Ans. (c) : $\because A^3 - B^3 = (A - B)(A^2 + B^2 + AB)$

$$27a^3 - 2\sqrt{2}b^3 = (3a)^3 - (\sqrt{2}b)^3$$

$$= (3a - \sqrt{2}b)(9a^2 + 2b^2 + 3\sqrt{2}ab)$$

192. If $x^4 + x^2y^2 + y^4 = 21$ and $x^2 + xy + y^2 = 7$, then

the value of $\left(\frac{1}{x^2} + \frac{1}{y^2}\right)$ is?

(a) $\frac{7}{4}$ (b) $\frac{5}{4}$

(c) $\frac{7}{3}$ (d) $\frac{5}{2}$

SSC CGL (Tier-I)-2019 – 03/03/2020 (Shift-II)

Ans. (b) : $x^2 + xy + y^2 = 7$ (1)

$$\therefore (x^2 - xy + y^2)(x^2 + xy + y^2) = x^4 + x^2y^2 + y^4$$

$$x^2 - xy + y^2 = \frac{x^4 + x^2y^2 + y^4}{x^2 + xy + y^2}$$

$$= \frac{21}{7} = 3 \dots \dots \dots (2)$$

For equation (1) + (2),

$$2(x^2 + y^2) = 10$$

$$x^2 + y^2 = 5$$

For equation (1) - (2),

$$2xy = 4$$

$$xy = 2$$

$$\therefore \frac{1}{x^2} + \frac{1}{y^2} = \frac{x^2 + y^2}{x^2y^2} = \frac{5}{4}$$

193. If $x - y = 4$ and $xy = 45$, then the value of $x^3 - y^3$ is:

- (a) 82 (b) 604
(c) 151 (d) 822

SSC CGL (Tier-I)-2019 – 03/03/2020 (Shift-I)

Ans. (b) : $\because x^3 - y^3 = (x - y)^3 + 3xy(x - y)$

$$= 64 + 3 \times 45 \times 4$$

$$= 64 + 540 = 604$$

194. If $2x^2 + y^2 + 8z^2 - 2\sqrt{2}xy + 4\sqrt{2}yz - 8zx = (Ax + y + Bz)^2$, then the value of $(A^2 + B^2 - AB)$ is:

- (a) 16 (b) 6
(c) 18 (d) 14

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-III)

Ans. (d) :

$$2x^2 + y^2 + 8z^2 - 2\sqrt{2}xy + 4\sqrt{2}yz - 8zx = (Ax + y + Bz)^2$$

$$\left(-\sqrt{2}x + y + 2\sqrt{2}z\right)^2 = (Ax + y + Bz)^2$$

On comparing the coefficients,

$$A = -\sqrt{2}, \quad B = 2\sqrt{2}$$

$$\therefore A^2 + B^2 - AB = 2 + 8 + 4 = 14$$

195. If $12x^2 - 21x + 1 = 0$, then what is the value of $9x^2 + (16x^2)^{-1}$?

- (a) $\frac{465}{16}$ (b) $\frac{429}{8}$
 (c) $\frac{417}{16}$ (d) $\frac{453}{8}$

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-III)

Ans. (c) : $12x^2 - 21x + 1 = 0$

$$12x + \frac{1}{x} = 21$$

$$3x + \frac{1}{4x} = \frac{21}{4}$$

On squaring both sides,

$$9x^2 + \frac{1}{16x^2} + 2 \times 3x \times \frac{1}{4x} = \frac{441}{16}$$

$$9x^2 + (16x^2)^{-1} = \frac{441}{16} - \frac{3}{2} = \frac{417}{16}$$

196. If $30x^2 - 15x + 1 = 0$, then what is the value of $25x^2 + (36x^2)^{-1}$?

- (a) $6\frac{1}{4}$ (b) $\frac{65}{12}$
 (c) $\frac{9}{2}$ (d) $\frac{55}{12}$

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-II)

Ans. (d) : $30x^2 - 15x + 1 = 0$

$$30x + \frac{1}{x} = 15$$

Dividing by 6

$$5x + \frac{1}{6x} = \frac{15}{6} = \frac{5}{2}$$

On squaring both sides,

$$25x^2 + \frac{1}{36x^2} + 2 \times 5x \times \frac{1}{6x} = \frac{25}{4}$$

$$25x^2 + \frac{1}{36x^2} = \frac{25}{4} - \frac{5}{3} = \frac{55}{12}$$

197. If $a + b + c = 7$ and $ab + bc + ca = -6$, then the value of $a^3 + b^3 + c^3 - 3abc$ is:

- (a) 463 (b) 469
 (c) 479 (d) 472

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-II)

Ans. (b) : $a^2 + b^2 + c^2 = (a+b+c)^2 - 2(ab+bc+ca)$
 $= 49 + 12 = 61$

$a^3 + b^3 + c^3 - 3abc = (a+b+c)(a^2+b^2+c^2-ab-bc-ca)$
 $= 7(61+6)$
 $= 7 \times 67 = 469$

198. If $P = \frac{x^4 - 8x}{x^3 - x^2 - 2x}$, $Q = \frac{x^2 + 2x + 1}{x^2 - 4x - 5}$ and $R = \frac{2x^2 + 4x + 8}{x - 5}$, then $(P \times Q) \div R$ is equal to:

- (a) $\frac{1}{2}$ (b) 2
 (c) 1 (d) 4

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-II)

Ans. (a) :

$$P = \frac{x^4 - 8x}{x^3 - x^2 - 2x} = \frac{x^3 - 8}{x^2 - x - 2} = \frac{(x-2)(x^2 + 2x + 4)}{(x-2)(x+1)}$$

$$= \frac{(x^2 + 2x + 4)}{(x+1)}$$

$$Q = \frac{x^2 + 2x + 1}{x^2 - 4x - 5} = \frac{(x+1)^2}{(x-5)(x+1)} = \frac{x+1}{x-5}$$

$$R = \frac{2(x^2 + 2x + 4)}{x - 5}$$

$$(P \times Q) \div R = \frac{x^2 + 2x + 4}{x+1} \times \frac{x+1}{x-5} \times \frac{x-5}{2(x^2 + 2x + 4)}$$

$$= \frac{1}{2}$$

199. If $5x + \frac{1}{3x} = 4$, then what is the value of $9x^2 + \frac{1}{25x^2}$?

- (a) $\frac{119}{25}$ (b) $\frac{174}{125}$
 (c) $\frac{144}{125}$ (d) $\frac{114}{25}$

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-I)

Ans. (d) :

$$5x + \frac{1}{3x} = 4$$

On multiplying by $\frac{3}{5}$

$$3x + \frac{1}{5x} = \frac{12}{5}$$

On squaring both sides

$$9x^2 + \frac{1}{25x^2} + 2 \times 3x \times \frac{1}{5x} = \frac{144}{25}$$

$$9x^2 + \frac{1}{25x^2} = \frac{144}{25} - \frac{6}{5} = \frac{114}{25}$$

200. If $a + b + c = 11$, $ab + bc + ca = 3$ and $abc = -135$, then what is the value of $a^3 + b^3 + c^3$?

- (a) 827 (b) 823
 (c) 925 (d) 929

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-I)

Ans. (a) : $\therefore a^3 + b^3 + c^3 - 3abc = (a+b+c)[(a+b+c)^2 - 3(ab+bc+ca)]$

$$a^3 + b^3 + c^3 + 405 = 11 [121 - 9]$$

$$a^3 + b^3 + c^3 = 1232 - 405 = 827$$

201. On simplification,

$$\frac{x^3 - y^3}{x[(x+y)^2 - 3xy]} \div \frac{y[(x-y)^2 + 3xy]}{x^3 + y^3} \times \frac{(x+y)^2 - (x-y)^2}{x^2 - y^2}$$

is equal to:

- (a) $\frac{1}{4}$ (b) 1
 (c) 4 (d) $\frac{1}{2}$

SSC CGL (Tier-I)-2019 – 04/03/2020 (Shift-I)

Ans. (c) :

$$\frac{x^3 - y^3}{x[(x+y)^2 - 3xy]} \div \frac{y[(x-y)^2 + 3xy]}{x^3 + y^3} \times \frac{(x+y)^2 - (x-y)^2}{x^2 - y^2}$$

$$= \frac{(x-y)(x^2 + xy + y^2)}{x(x^2 + y^2 - xy)} \times \frac{(x+y)(x^2 - xy + y^2)}{y(x^2 + y^2 + xy)} \times \frac{4xy}{(x+y)(x-y)}$$

$$= 4$$

202. If $x^4 + x^2y^2 + y^4 = 273$ and $x^2 - xy + y^2 = 13$, then the value of xy is:

- (a) 6 (b) 10
 (c) 8 (d) 4

SSC CGL (Tier-I)-2019 – 05/03/2020 (Shift-II)

Ans. (d) : $x^2 + xy + y^2 = \frac{x^4 + x^2y^2 + y^4}{x^2 - xy + y^2}$

$$x^2 + xy + y^2 = \frac{273}{13}$$

$$x^2 + xy + y^2 = 21 \dots\dots\dots(i)$$

$$x^2 - xy + y^2 = 13 \dots\dots\dots(ii)$$

From equation (i) and equation (ii),

$$2xy = 8$$

$$xy = 4$$

203. If $20x^2 - 30x + 1 = 0$, then what is the value of

$$25x^2 + \frac{1}{16x^2} ?$$

- (a) $53\frac{1}{2}$ (b) $58\frac{3}{4}$
 (c) $58\frac{1}{2}$ (d) $53\frac{3}{4}$

SSC CGL (Tier-I)-2019 – 05/03/2020 (Shift-II)

Ans. (d) : $20x^2 - 30x + 1 = 0$

$$20x + \frac{1}{x} = 30$$

$$5x + \frac{1}{4x} = \frac{15}{2}$$

On squaring both sides,

$$25x^2 + \frac{1}{16x^2} + 2 \times 5x \times \frac{1}{4x} = \frac{225}{4}$$

$$25x^2 + \frac{1}{16x^2} = \frac{225}{4} - \frac{5}{2} = \frac{215}{4} = 53\frac{3}{4}$$

204. If $16a^4 + 36a^2b^2 + 81b^4 = 91$ and $4a^2 + 9b^2 - 6ab = 13$, then what is the value of $3ab$?

- (a) $-\frac{3}{2}$ (b) $\frac{3}{2}$
 (c) 5 (d) -3

SSC CGL (Tier-I)-2019 – 05/03/2020 (Shift-I)

Ans. (a) :

$$16a^4 + 36a^2b^2 + 81b^4 = (4a^2 + 9b^2 - 6ab)^2 \quad (4a^2 + 9b^2 + 6ab)$$

From formula

$$4a^2 + 9b^2 + 6ab = \frac{91}{13} = 7 \dots\dots\dots(i)$$

$$4a^2 + 9b^2 - 6ab = 13 \dots\dots\dots(ii)$$

From equation (i) – equation (ii)

$$12ab = -6$$

$$3ab = \frac{-6}{4} = \frac{-3}{2}$$

205. If $x^2 - 2\sqrt{5}x + 1 = 0$, then what is the value of $x^5 + \frac{1}{x^5}$?

- (a) $408\sqrt{5}$ (b) $612\sqrt{5}$
 (c) $406\sqrt{5}$ (d) $610\sqrt{5}$

SSC CGL (Tier-I)-2019 – 05/03/2020 (Shift-I)

Ans. (d) : $x^2 - 2\sqrt{5}x + 1 = 0$

$$x + \frac{1}{x} = 2\sqrt{5}$$

$$\therefore x^5 + \frac{1}{x^5} = \left(x^2 + \frac{1}{x^2}\right)\left(x^3 + \frac{1}{x^3}\right) - \left(x + \frac{1}{x}\right)$$

$$\text{then, } x^2 + \frac{1}{x^2} = (2\sqrt{5})^2 - 2 = 18$$

$$x^3 + \frac{1}{x^3} = (2\sqrt{5})^3 - 3 \times 2\sqrt{5} = 34\sqrt{5}$$

$$\therefore \left(x^2 + \frac{1}{x^2}\right)\left(x^3 + \frac{1}{x^3}\right) = 18 \times 34\sqrt{5}$$

$$\therefore x^5 + \frac{1}{x^5} = 612\sqrt{5} - 2\sqrt{5} = 610\sqrt{5}$$

206. Find the product of $(a+b+2c)(a^2+b^2+4c^2-ab-2bc-2ca)$.

- (a) $a^3 + b^3 + 8c^3 - 6abc$ (b) $a^3 + b^3 + 8c^3 - 2abc$
 (c) $a^3 + b^3 + 8c^3 - abc$ (d) $a^3 + b^3 + 6c^3 - 6abc$

SSC CGL (Tier-I)-2019 – 07/03/2020 (Shift-III)

Ans. (a) : $(a+b+2c)(a^2+b^2+4c^2-ab-2bc-2ca)$

$$= (a+b+2c)[a^2+b^2+(2c)^2-ab-2bc-2ca]$$

$$\therefore a^3 + b^3 + (2c)^3 - 3 \times a \times b \times (2c)$$

$$= a^3 + b^3 + 8c^3 - 6abc$$

207. If $a^4 + \frac{1}{a^4} = 50$, $a > 0$, then find the value of

$$a^3 + \frac{1}{a^3}.$$

(a) $\sqrt{2(1-\sqrt{13})}(-1+2\sqrt{13})$

(b) $\sqrt{2(1+\sqrt{13})}(-1-2\sqrt{13})$

(c) $\sqrt{2(1+\sqrt{13})}(-1+2\sqrt{13})$

(d) $\sqrt{2(1+\sqrt{13})}+(-1+2\sqrt{13})$

SSC CGL (Tier-I)-2019 – 07/03/2020 (Shift-III)

Ans. (c) : $a^4 + \frac{1}{a^4} = 50$

$$a^2 + \frac{1}{a^2} = \sqrt{52}$$

$$\left(a + \frac{1}{a}\right)^2 = 2\sqrt{13} + 2$$

$$a + \frac{1}{a} = \sqrt{2(1 + \sqrt{13})}$$

if $a + \frac{1}{a} = k$ and $a^3 + \frac{1}{a^3} = k^3 - 3k$

$$\begin{aligned} \therefore a^3 + \frac{1}{a^3} &= 2(1 + \sqrt{13})\sqrt{2(1 + \sqrt{13})} - 3\sqrt{2(1 + \sqrt{13})} \\ &= \sqrt{2(1 + \sqrt{13})}(-1 + 2\sqrt{13}) \end{aligned}$$

208. If $a^2 + b^2 + c^2 = 300$ and $ab + bc + ca = 50$, then what is the value of $a + b + c$? (Given that a, b and c are all positive).

- (a) 15 (b) 20
(c) 22 (d) 25

SSC CGL (Tier-I)-2019 – 07/03/2020 (Shift-II)

Ans. (b) : $(a+b+c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$
 $= 300 + 100 = 400$
 $a + b + c = 20$

209. If $1 - 64x^3 - 12x + px^2 = (1-4x)^3$, then the value of p is:

- (a) 48 (b) -12
(c) -48 (d) 16

SSC CGL (Tier-I)-2019 – 07/03/2020 (Shift-I)

Ans. (a) : $1 - 64x^3 - 12x + px^2 = (1-4x)^3$
 $1 - 64x^3 - 12x + px^2 = 1 - 64x^3 - 12x + 48x^2$
 On comparing both sides,
 $p = 48$

210. If $a^2 + b^2 - c^2 = 0$, then the value of $\frac{2(a^6 + b^6 - c^6)}{3a^2b^2c^2}$ is:

- (a) 1 (b) 0
(c) 2 (d) 3

SSC CGL (Tier-I)-2019 – 09/03/2020 (Shift-II)

Ans. (*) : $\because a^2 + b^2 - c^2 = 0$
 $a^2 + b^2 = c^2$ (i)
 On cubing both sides,
 $(a^2 + b^2)^3 = c^6$
 $a^6 + b^6 + 3a^2b^2(a^2 + b^2) = c^6$
 $a^6 + b^6 - c^6 = -3a^2b^2c^2$ -----($\because a^2 + b^2 = c$)
 $\frac{a^6 + b^6 - c^6}{3a^2b^2c^2} = -1$
 On multiplying by 2 of both sides
 $\frac{2(a^6 + b^6 - c^6)}{3a^2b^2c^2} = -2$

Note- SSC (Staff selection commission) has considered the answer to this question as 2, while the correct answer would be -2.

211. Expand : $(4a + 3b + 2c)^2$

- (a) $4a^2 + 3b^2 + 2c^2 + 24ab + 12bc + 16ca$
 (b) $16a^2 - 9b^2 + 4c^2 - 24ab + 12bc - 16ca$
 (c) $16a^2 + 9b^2 + 4c^2 + 24ab + 12bc + 16ca$
 (d) $16a^2 + 9b^2 + 4c^2 - 24ab - 12bc - 16ca$

SSC CGL (Tier-I)-2019 – 09/03/2020 (Shift-I)

Ans. (c) : We know that formula:-

$$\begin{aligned} \therefore (a + b + c)^2 &= (a^2 + b^2 + c^2 + 2ab + 2bc + 2ca) \\ (4a + 3b + 2c)^2 &= 16a^2 + 9b^2 + 4c^2 + 24ab + 12bc + 16ca \end{aligned}$$

212. If $x + y = 10$ and $xy = 4$, then what is the value of $x^4 + y^4$?

- (a) 8464 (b) 8432
(c) 7478 (d) 6218

SSC CGL (Tier-II) 21-02-2018

Ans. (b) : Given-

$x + y = 10$ (i)
 $xy = 4$ (ii)
 $\therefore x^4 + y^4 = (x^2 + y^2)^2 - 2x^2y^2$
 $\therefore x^4 + y^4 = [(x+y)^2 - 2xy]^2 - 2(xy)^2$ (iii)

By putting the value from equation (i) and (ii) in equation (iii)

$$\begin{aligned} x^4 + y^4 &= [(10)^2 - 2 \times 4]^2 - 2 \times (4)^2 \\ &= (92)^2 - 2 \times 16 \\ &= 8464 - 32 \\ &= \boxed{8432} \end{aligned}$$

213. If $a + b + c = 9$, $ab + bc + ca = 26$, $a^3 + b^3 = 91$, $b^3 + c^3 = 72$ and $c^3 + a^3 = 35$, then what is the value of abc ?

- (a) 48 (b) 24
(c) 36 (d) 42

SSC CGL (Tier-II) 21-02-2018

Ans. (b) : $a^3 + b^3 = 91$ -----(i)
 $b^3 + c^3 = 72$ -----(ii)
 $c^3 + a^3 = 35$ -----(iii)

By adding

$$\begin{aligned} 2(a^3 + b^3 + c^3) &= 198 \\ a^3 + b^3 + c^3 &= 99 \end{aligned}$$

$$\begin{aligned} \therefore a^3 + b^3 + c^3 - 3abc &= (a+b+c) [(a+b+c)^2 - 3(ab+bc+ca)] \\ 99 - 3abc &= 9(81 - 78) \\ 3abc &= 99 - 27 = 72 \\ abc &= 24 \end{aligned}$$

214. If $x^3 - 4x^2 + 19 = 6(x-1)$, then what is the value of $[x^2 + (1/x - 4)]^2$?

- (a) 3 (b) 5
(c) 6 (d) 8

SSC CGL (Tier-II) 21-02-2018

Ans. (c) : Given-

$$\begin{aligned} x^3 - 4x^2 + 19 &= 6(x-1) \\ \Rightarrow x^3 - 4x^2 &= 6x - 6 - 19 \\ \Rightarrow x^3 - 4x^2 &= 6x - 25 \text{ (i)} \end{aligned}$$

$$\therefore x^2 + \frac{1}{(x-4)} = \frac{x^3 - 4x^2 + 1}{x-4}$$

$$= \frac{6x-25+1}{x-4} \quad (\text{From equation (i)})$$

$$= \frac{6(x-4)}{(x-4)}$$

$$= \boxed{6}$$

215. If $x + y + z = 22$ and $xy + yz + zx = 35$, then what is the value of $(x-y)^2 + (y-z)^2 + (z-x)^2$?
- (a) 793 (b) 681
(c) 758 (d) 715

SSC CGL (Tier-II) 20-02-2018

Ans. (c) $(x-y)^2 + (y-z)^2 + (z-x)^2$
 $= [x^2+y^2-2xy+y^2+z^2-2yz+z^2+x^2-2zx]$
 $= 2(x^2+y^2+z^2-xy-yz-zx)$
 $= 2[(x+y+z)^2 - 3(xy+yz+zx)]$
 $= 2[(22)^2 - 3 \times 35]$
 $= 2[484 - 105]$
 $= 2 \times 379$
 $= 758$

216. If α and β are the roots of the equation $x^2+x-1=0$, then what is the equation whose roots are α^5 and β^5 ?
- (a) $x^2+7x-1=0$ (b) $x^2-7x-1=0$
(c) $x^2-11x-1=0$ (d) $x^2+11x-1=0$

SSC CGL (Tier-II) 19-02-2018

Ans. (d) : $x^2 + x - 1 = 0$
 $\alpha + \beta = -1$
 $\alpha \cdot \beta = -1$
 $\alpha^2 + \beta^2 = (-1)^2 + 2 = 3$
 $\alpha^3 + \beta^3 = (-1)^3 - 3 \times (-1) \times (-1) = -1 - 3 = -4$
 $(\alpha^2 + \beta^2)(\alpha^3 + \beta^3) = \alpha^5 + \alpha^2\beta^2 + \beta^2\alpha^3 + \beta^5$
 $3 \times (-4) = \alpha^5 + \beta^5 + \alpha^2\beta^2(\alpha + \beta)$
 $-12 = \alpha^5 + \beta^5 + 1 \times (-1)$
 $\alpha^5 + \beta^5 = -11$
 \therefore Quadratic equation
 $x^2 - (\alpha^5 + \beta^5)x + (\alpha\beta)^5 = 0$
 $x^2 + 11x - 1 = 0$

217. If $x + (1/x) = (\sqrt{3}+1)/2$, then what is the value of $x^4 + (1/x^4)$?
- (a) $(4\sqrt{3}-1)/4$ (b) $(4\sqrt{3}+1)/2$
(c) $(-4\sqrt{3}-1)/4$ (d) $(-4\sqrt{3}-1)/2$

SSC CGL (Tier-II) 19-02-2018

Ans. (c) : $x + \frac{1}{x} = \frac{\sqrt{3}+1}{2}$ (Squaring in both sides)

$$x^2 + \frac{1}{x^2} = \left(\frac{\sqrt{3}+1}{2}\right)^2 - 2$$

$$= \frac{4+2\sqrt{3}}{4} - 2 = \frac{2\sqrt{3}-4}{4} = \frac{\sqrt{3}-2}{2}$$

$$x^4 + \frac{1}{x^4} = \left(\frac{\sqrt{3}-2}{2}\right)^2 - 2$$

$$= \frac{7-4\sqrt{3}}{4} - 2 = \frac{-4\sqrt{3}-1}{4}$$

218. $A = (x^8-1)/(x^4+1)$ and $B = (y^4-1)/(y^2+1)$. If $x = 2$ and $y = 9$, then what is the value of $A^2 + 2AB + AB^2$?
- (a) 96475 (b) 98625
(c) 92425 (d) 89125

SSC CGL (Tier-II) 18-02-2018

Ans. (b) : Given-

$$A = \frac{x^8-1}{x^4+1} \quad \& \quad B = \frac{y^4-1}{y^2+1}$$

From formula

$$(a-b)(a+b) = a^2 - b^2$$

By solving the equation A and B

$$A = \frac{(x^4-1)(x^4+1)}{(x^4+1)} \quad \& \quad B = \frac{(y^2-1)(y^2+1)}{(y^2+1)}$$

$$A = (x^4-1) \quad \& \quad B = (y^2-1)$$

$$\text{If } x = 2, \quad \& \quad y = 9$$

Then-

$$A = 2^4 - 1 = 15$$

$$B = 9^2 - 1 = 80$$

By putting the value of A and B

$$\Rightarrow A^2 + AB^2 + 2AB$$

$$\Rightarrow (15)^2 + 15 \times (80)^2 + 2 \times 15 \times 80$$

$$= 225 + 96000 + 2400 = 98625$$

219. If $x = (\sqrt{5}+1)$ and $y = (\sqrt{5}-1)$ then what is the value of $(x^2/y^2) + (y^2/x^2) + 4\left(\frac{x}{y} + \frac{y}{x}\right) + 6$?

- (a) 31 (b) $23\sqrt{5}$
(c) $27\sqrt{5}$ (d) 25

SSC CGL (Tier-II) 18-02-2018

Ans. (d) : Given,

$$x = \sqrt{5} + 1 \dots \dots \dots (1)$$

$$y = \sqrt{5} - 1 \dots \dots \dots (2)$$

From equation (1) and (2)

$$x + y = 2\sqrt{5} \quad \& \quad xy = 4$$

$$x^2 + y^2 = (x+y)^2 - 2xy$$

$$= 20 - 2 \times 4$$

$$x^2 + y^2 = 12$$

$$= \frac{x^2}{y^2} + \frac{y^2}{x^2} + 4 \left[\left(\frac{x}{y}\right) + \left(\frac{y}{x}\right) \right] + 6$$

$$= \frac{x^4 + y^4}{x^2y^2} + 4 \left[\frac{x^2 + y^2}{xy} \right] + 6$$

$$= \frac{(x^2 + y^2)^2 - 2x^2y^2}{x^2y^2} + 4 \left[\frac{(x+y)^2 - 2xy}{xy} \right] + 6 \dots \dots \dots (3)$$

By putting the value in equation (3)

$$\Rightarrow \frac{(12)^2 - 2 \times 16}{16} + \frac{4 \times (12)}{4} + 6$$

$$= 7 + 18 = 25$$

220. If $x = 2 + \sqrt{3}$, $y = 2 - \sqrt{3}$ and $z = 1$, then what is the value of $(x/yz) + (y/xz) + (z/xy) + 2 [(1/x) + (1/y) + (1/z)]$?

- (a) 25 (b) 22
(c) 17 (d) 43

SSC CGL (Tier-II) 18-02-2018

Ans. (a) : Given-

$$x = 2 + \sqrt{3}, y = 2 - \sqrt{3}, z = 1$$

$$x \times y \times z = (2 + \sqrt{3}) \times (2 - \sqrt{3}) \times 1 = 1$$

$$x + y + z = 2 + \sqrt{3} + 2 - \sqrt{3} + 1 = 5$$

$$x^2 + y^2 + z^2 = (2 + \sqrt{3})^2 + (2 - \sqrt{3})^2 + 1$$

$$7 + 4\sqrt{3} + 7 - 4\sqrt{3} + 1$$

$$= 7 + 7 + 1$$

$$= 15$$

We know that

$$(x+y+z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)$$

$$5^2 = 15 + 2(xy + yz + zx)$$

$$xy + yz + zx = \frac{10}{2} = 5$$

$$\frac{x}{yz} + \frac{y}{xz} + \frac{z}{xy} + 2 \left[\frac{1}{x} + \frac{1}{y} + \frac{1}{z} \right]$$

$$\Rightarrow \frac{x^2 + y^2 + z^2}{xyz} + 2 \left[\frac{xy + yz + zx}{xyz} \right] \text{ (Putting the value)}$$

$$\frac{15}{1} + 2 \left[\frac{5}{1} \right] \Rightarrow 15 + 10 = 25$$

221. If $f(x) = (x-2)/(x^2 + Px + 4)$ and $(x-3)$ is a factor of $f(x)$, then what is the value of P ?

- (a) 4 (b) -4
(c) -13/3 (d) 13/3

SSC CGL (Tier-II) 9-3-2018

Ans. (c) : $f(x) = (x-2)/(x^2 + Px + 4)$

$\therefore (x-3)$, is a factor of $f(x)$

Then putting the value of $x = 3$

Remainder = 0

$$(3-2)(9 + 3P + 4) = 0$$

$$3P = -13 \Rightarrow P = \frac{-13}{3}$$

222. If $[x - (1/x)] = 2$, then what is the value of $[x^6 - (1/x^6)]$?

- (a) $114\sqrt{3} + 1$ (b) $134\sqrt{2}$
(c) $142\sqrt{2} + 3$ (d) $140\sqrt{2}$

SSC CGL (Tier-II) 9-3-2018

Ans. (d) :

$$x - \frac{1}{x} = 2 \quad \text{(squaring both sides)}$$

$$x^3 - \frac{1}{x^3} = (2)^3 + 3 \times 2 = 14 \dots \dots (1)$$

$$\text{Again } x + \frac{1}{x} = \sqrt{(2)^2 + 4} = \sqrt{8}$$

$$x^3 + \frac{1}{x^3} = (\sqrt{8})^3 - 3\sqrt{8} = 5\sqrt{8} \dots \dots (2)$$

By multiplying the equation (i) and (ii)

$$x^6 - \frac{1}{x^6} = 70\sqrt{8} = 140\sqrt{2}$$

223. x, y and z are real numbers. If $x^3 + y^3 + z^3 = 13$, $x + y + z = 1$ and $xyz = 1$, then what is the value of $xy + yz + zx$?

- (a) -1 (b) 1
(c) 3 (d) -3

SSC CGL (Tier-II) 17-2-2018

Ans. (d)

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)$$

$$13 - 3 \times 1 = 1 \left((x + y + z)^2 - 3(xy + yz + zx) \right)$$

$$10 = 1^2 - 3(xy + yz + zx)$$

$$3(xy + yz + zx) = 1 - 10 = -9$$

$$xy + yz + zx = -3$$

224. If $x^3 + y^3 + z^3 = 3(1 + xyz)$, $P = y + z - x$, $Q = z + x - y$ and $R = x + y - z$, then what is the value of $P^3 + Q^3 + R^3 - 3PQR$?

- (a) 9 (b) 8
(c) 12 (d) 6

SSC CGL (Tier-II) 17-2-2018

Ans. (c) : From value putting,

$$y = z = 0$$

$$x^3 + 0 = 3(1 + 0)$$

$$x^3 = 3$$

$$P = 0 + 0 - x = -x$$

$$Q = 0 + x - 0 = x$$

$$R = x + 0 - 0 = x$$

$$P^3 + Q^3 + R^3 - 3PQR = (-x)^3 + x^3 + x^3 - 3 \times (-x) \times x \times x$$

$$= x^3 + 3x^3$$

$$= 4x^3$$

$$= 4 \times 3 = 12$$

225. The value of $\frac{(4.6)^4 + (5.4)^4 + (24.84)^2}{(4.6)^2 + (5.4)^2 + 24.84}$ is :

- (a) 24.42 (b) 25.48
(c) 24.24 (d) 25.42

SSC CGL (Tier-II) 13-09-2019

Ans. (b) :

$$\frac{(4.6)^4 + (5.4)^4 + (24.84)^2}{(4.6)^2 + (5.4)^2 + 24.84}$$

$$\text{Hence } x = 4.6, y = 5.4$$

$$x^4 + y^4 + x^2y^2 = (x^2 + y^2 + xy)(x^2 + y^2 - xy)$$

$$= \frac{[(4.6)^2 + (5.4)^2 + 4.6 \times 5.4][(4.6)^2 + (5.4)^2 - 4.6 \times 5.4]}{(4.6)^2 + (5.4)^2 + 4.6 \times 5.4}$$

$$= (4.6 + 5.4)^2 - 3 \times 4.6 \times 5.4$$

$$= 100 - 74.52 = 25.48$$

226. Let $x = \sqrt[6]{27} - \sqrt{6\frac{3}{4}}$ and $y = \frac{\sqrt{45} + \sqrt{605} + \sqrt{245}}{\sqrt{80} + \sqrt{125}}$, then the value of $x^2 + y^2$ is?

(a) $\frac{223}{36}$ (b) $\frac{221}{36}$
(c) $\frac{221}{9}$ (d) $\frac{227}{9}$

SSC CGL (Tier-II) 13-09-2019

Ans. (a) :

$$x = \sqrt[6]{27} - \sqrt{6\frac{3}{4}}$$

$$= (3^3)^{\frac{1}{6}} - \sqrt{\frac{27}{4}}$$

$$= \sqrt{3} - \frac{3\sqrt{3}}{2} = \frac{-\sqrt{3}}{2}$$

$$y = \frac{\sqrt{45} + \sqrt{605} + \sqrt{245}}{\sqrt{80} + \sqrt{125}} = \frac{3\sqrt{5} + 11\sqrt{5} + 7\sqrt{5}}{4\sqrt{5} + 5\sqrt{5}} = \frac{7}{3}$$

$$x^2 + y^2 = \frac{3}{4} + \frac{49}{9} = \frac{27 + 196}{36} = \frac{223}{36}$$

227. If $8x^3 - 27y^3 = (Ax + By)(Cx^2 - Dy^2 + 6xy)$, then $(A + B + C - D)$ is equal to :

(a) -12 (b) 12
(c) 9 (d) 15

SSC CGL (Tier-II) 12-09-2019

Ans. (b) : Given,

$$8x^3 - 27y^3 = (Ax + By)(Cx^2 - Dy^2 + 6xy)$$

$$(2x - 3y)(4x^2 + 9y^2 + 6xy) = (Ax + By)(Cx^2 - Dy^2 + 6xy)$$

On comparing both sides,
 $A = 2, C = 4$
 $B = -3, D = -9$
 $\therefore (A+B+C-D) = 2 - 3 + 4 + 9 = 12$

228. If $x = \frac{\sqrt{5} - \sqrt{3}}{\sqrt{5} + \sqrt{3}}$ and y is the reciprocal of x , then what is the value of $(x^3 + y^3)$?

(a) 504 (b) 476
(c) 472 (d) 488

SSC CGL (Tier-II) 12-09-2019

Ans. (d) :

$$\therefore xy = 1$$

$$x + y = \frac{\sqrt{5} - \sqrt{3}}{\sqrt{5} + \sqrt{3}} + \frac{\sqrt{5} + \sqrt{3}}{\sqrt{5} - \sqrt{3}}$$

$$= \frac{5 + 3 - 2\sqrt{15} + 5 + 3 + 2\sqrt{15}}{2} = 8$$

$$x^3 + y^3 = (x+y)^3 - 3xy(x+y)$$

$$= (x+y)[(x+y)^2 - 3xy]$$

$$= 8[64 - 3]$$

$$= 8 \times 61 = 488$$

229. If $x^4 - 83x^2 + 1 = 0$, then value of $x^3 - x^{-3}$ can be:

- (a) 758 (b) 739
(c) 737 (d) 756

SSC CGL (Tier-II) 12-09-2019

Ans. (d) : $x^4 - 83x^2 + 1 = 0$

$$x^4 + 1 = 83x^2 \Rightarrow x^2 + \frac{1}{x^2} = 83$$

$$\left(x - \frac{1}{x}\right)^2 = 81 \Rightarrow \left(x - \frac{1}{x}\right) = 9$$

\therefore On cubing both sides,

$$x^3 - \frac{1}{x^3} = 729 + 3 \times 9 = 729 + 27$$

$\therefore x^3 - x^{-3} = 756$

230. If $x + y + z = 2$, $xy + yz + zx = -11$ and $xyz = -12$, then what is the value of $\sqrt{x^3 + y^3 + z^3 - 2}$?

- (a) 12 (b) 9
(c) 6 (d) 8

SSC CGL (Tier-II) 12-09-2019

Ans. (c) : Value putting,

$$\therefore xyz = -12 = 1 \times (-3) \times 4$$

Taking the value $x = 1, y = -3, z = 4$

$$x + y + z = 1 - 3 + 4 = 2$$

$$xy + yz + zx = -3 - 12 + 4 = -11$$

$$\therefore \sqrt{x^3 + y^3 + z^3 - 2} = \sqrt{1^3 + (-3)^3 + 4^3 - 2} = \sqrt{1 - 27 + 64 - 2} = \sqrt{63 - 27} = \sqrt{36} = 6$$

231. If $x + \frac{1}{16x} = 3$, then the value of

$$16x^3 + \frac{1}{256x^3} \text{ is:}$$

- (a) 423 (b) 441
(c) 414 (d) 432

SSC CGL (Tier-II) 12-09-2019

Ans. (a) :

$$x + \frac{1}{16x} = 3$$

On cubing both sides,

$$x^3 + \frac{1}{4096x^3} + 3 \times x \times \frac{1}{16x} \left(x + \frac{1}{16x}\right) = 27$$

$$x^3 + \frac{1}{4096x^3} + \frac{3}{16} \times 3 = 27$$

$$x^3 + \frac{1}{4096x^3} = 27 - \frac{9}{16} = \frac{432 - 9}{16} = \frac{423}{16}$$

Multiplying by 16 in both sides,

$$16x^3 + \frac{1}{256x^3} = 16 \times \frac{423}{16} = 423$$